




ELSEVIER



British Mycological  
Society promoting fungal science

journal homepage: [www.elsevier.com/locate/mycres](http://www.elsevier.com/locate/mycres)

## Review

## Medicinal importance of fungal $\beta$ -(1 $\rightarrow$ 3), (1 $\rightarrow$ 6)-glucans

Jiezhong CHEN<sup>a,\*</sup>, Robert SEVIOUR<sup>b</sup>

<sup>a</sup>Cancer Biology Program, Diamantina Institute for Cancer, Immunology and Metabolic Medicine, University of Queensland, Brisbane, Queensland 4102, Australia

<sup>b</sup>Biotechnology Research Centre, School of Pharmacy and Applied Science, Faculty of Science, Technology and Engineering, La Trobe University Bendigo, Victoria 3550, Australia

## ARTICLE INFO

## Article history:

Received 18 September 2006

Received in revised form

25 January 2007

Accepted 19 February 2007

Published online 7 March 2007

Corresponding Editor:

Mark Ramsdale

## Keywords:

Anticancer

 $\beta$ -glucans

Dectin-1

Fungi

Immunomodulators

## ABSTRACT

Non-cellulosic  $\beta$ -glucans are now recognized as potent immunological activators, and some are used clinically in China and Japan. These  $\beta$ -glucans consist of a backbone of glucose residues linked by  $\beta$ -(1  $\rightarrow$  3)-glycosidic bonds, often with attached side-chain glucose residues joined by  $\beta$ -(1  $\rightarrow$  6) linkages. The frequency of branching varies. The literature suggests  $\beta$ -glucans are effective in treating diseases like cancer, a range of microbial infections, hypercholesterolaemia, and diabetes. Their mechanisms of action involve them being recognized as non-self molecules, so the immune system is stimulated by their presence. Several receptors have been identified, which include: dectin-1, located on macrophages, which mediates  $\beta$ -glucan activation of phagocytosis and production of cytokines, a response co-ordinated by the toll-like receptor-2. Activated complement receptors on natural killer cells, neutrophils, and lymphocytes, may also be associated with tumour cytotoxicity. Two other receptors, scavenger and lactosylceramide, bind  $\beta$ -glucans and mediate a series of signal pathways leading to immunological activation. Structurally different  $\beta$ -glucans appear to have different affinities toward these receptors and thus generate markedly different host responses. However, the published data are not always easy to interpret as many of the earlier studies used crude  $\beta$ -glucan preparations with, for the most part, unknown chemical structures. Careful choice of  $\beta$ -glucan products is essential if their benefits are to be optimized, and a better understanding of how  $\beta$ -glucans bind to receptors should enable more efficient use of their biological activities.

© 2007 The British Mycological Society. Published by Elsevier Ltd. All rights reserved.

### Introduction

Non-cellulosic  $\beta$ -glucans are recognized as potent immunological stimulators in humans and some are now used clinically in China and Japan. These  $\beta$ -glucans appear to possess potential for treating several diseases. About half the mass of the fungal cell wall consists of  $\beta$ -glucans (Seviour *et al.*

1992; Klis *et al.* 2001; McIntosh *et al.* 2005), but many are also excreted into the growth medium, making their recovery, purification and chemical characterization much simpler (Seviour *et al.* 1992; Schmid *et al.* 2001). Their homeostasis is maintained by  $\beta$ -glucan synthases and  $\beta$ -(1  $\rightarrow$  3) or  $\beta$ -(1  $\rightarrow$  6)-glucanase activities (Pitson *et al.* 1996a, 1996b; Jayus *et al.* 2004; Martin *et al.* 2006). Inhibition of  $\beta$ -glucan synthases by

\* Corresponding author.

E-mail address: [jchen@cicr.uq.edu.au](mailto:jchen@cicr.uq.edu.au)

0953-7562/\$ - see front matter © 2007 The British Mycological Society. Published by Elsevier Ltd. All rights reserved.

doi:10.1016/j.mycres.2007.02.011

compounds like caspofungin, although expensive, has been successful in treating *Aspergillus* spp. and *Candida* spp. infections in humans (Walsh et al. 2004).

Non-cellulosic  $\beta$ -glucans consist of a backbone of glucose residues usually joined by  $\beta$ -(1 $\rightarrow$ 3) linkages, to which glucose side-chain residues are often attached, as shown in examples in Fig 1 (Schmid et al. 2001; McIntosh et al. 2005). In some  $\beta$ -glucans no side-chain substitution occurs, as with the bacterial  $\beta$ -glucan, curdlan, which contains only  $\beta$ -(1 $\rightarrow$ 3)-glucosidic linkages (McIntosh et al. 2005). No unbranched  $\beta$ -(1 $\rightarrow$ 3) or  $\beta$ -(1 $\rightarrow$ 6) fungal  $\beta$ -glucans are known, although the extent of the side-chain substitutions can vary considerably (Schmid et al. 2001). For example, many including schizophyllan (also named sizofiran) from *Schizophyllum commune*, and scleroglucan from *Sclerotium glaucum* both have a  $\beta$ -(1 $\rightarrow$ 3)-linked backbone, with on average one  $\beta$ -(1 $\rightarrow$ 6)-glucose substitution every three backbone residues (Johnson et al. 1963; Kikumoto & Kimura 1971). Epiglucan from *Epicoccum nigrum* also has a backbone of  $\beta$ -(1 $\rightarrow$ 3)-linked glucose residues, but now with two  $\beta$ -(1 $\rightarrow$ 6) substitutions on average every three residues, while the  $\beta$ -(1 $\rightarrow$ 3)-linked backbone of lentinan (also called SPG) from *Lentinus edodes* has two  $\beta$ -(1 $\rightarrow$ 6) side chains every five residues (Sasaki & Takasuka 1976; Schmid et al. 2001). Pestalotian produced extracellularly by *Pestalotia* sp. 815 has a  $\beta$ -(1 $\rightarrow$ 3)-linked backbone, but with three  $\beta$ -(1 $\rightarrow$ 6) side chains every five residues (Misaki et al. 1984). Many other fungal  $\beta$ -glucans have been described, but their detailed structures and branching frequencies are still mostly unclear (Seviour et al. 1992). It appears that to be effective these glucans must contain  $\beta$ -(1 $\rightarrow$ 3) or  $\beta$ -(1 $\rightarrow$ 6) linkages, but little else is known about how frequencies of branching and other chemical and physical properties determine their effectiveness, as discussed later (Demleitner et al. 1992). Examples of some non-cellulosic fungal  $\beta$ -glucans of possible medical importance are listed in Table 1.

Walls of pathogenic fungi also contain  $\beta$ -(1 $\rightarrow$ 3) or  $\beta$ -(1 $\rightarrow$ 6)-glucans (Taylor et al. 2007; Saijo et al. 2007). Although (for practical reasons) these are not applied clinically, they are probably important in eliciting anti-fungal host responses. However, which wall component is the stimulator/s is not always clear, and it is also not clear whether all fungi behave in the same mechanistic way. For example, *A. fumigatus* synthesizes a  $\beta$ -(1 $\rightarrow$ 3)-glucan with 4%  $\beta$ -(1 $\rightarrow$ 6) branches and a linear  $\beta$ -(1 $\rightarrow$ 3),(1 $\rightarrow$ 4)-glucan (Fontaine et al. 2000) and both stimulate immune responses in mice (Hohl et al. 2005). Yet the role of *Candida albicans* cell wall  $\beta$ -glucans appears uncertain. Evidence suggest some competition with laminarin (Netea et al. 2006), which, although a  $\beta$ -glucan and able preferentially to bind to the cell receptors, fails to induce any subsequent host immune responses. However, walls of *Pneumocystis carinii* f. sp. *muris* also contain a  $\beta$ -(1 $\rightarrow$ 3)-glucan, and digestion by a  $\beta$ -(1 $\rightarrow$ 3)-glucanase greatly reduces its immunostimulatory effect (Vassallo et al. 2000). In the living organism, wall  $\beta$ -glucans are apparently shielded by mannan layers, but  $\beta$ -glucans become exposed after heat treatment, and induce stronger host immunoresponses (Gantner et al. 2005). Further work with well-characterized  $\beta$ -glucans from the walls of other pathogenic fungi is needed to clarify their roles, if any, in host defence mechanisms.

Natural products containing fungal  $\beta$ -glucans have been consumed for probably thousands of years, and anecdotally, especially in China and Japan (Mayell 2001; Lindequist et al. 2005), basidiocarps have long been considered to improve general health.  $\beta$ -Glucans were only recently recognized as the effective ingredients (Lucas et al. 1958) (Williams & Di Luzio 1980). Subsequently, detailed investigations into their influence on health have been reported, mainly using animal models (Vetvicka & Yvin 2004). Several fungal  $\beta$ -glucans appear to be effective immunomodulators, and they appear to impact positively on cancers and several bacterial infections (Brown & Gordon 2001; Vetvicka & Yvin 2004). Although not yet medically prescribed in most countries, their administration is now increasing. For example, in Japan lentinan is approved for clinical treatment of gastric and colorectal cancers, and encouraging survival rates have been reported (Nakano et al. 1999; Munemoto et al. 2002). Some of their possible mechanisms of action have also been elucidated (Brown & Gordon 2001, 2005). In this review, we discuss the potential of fungal  $\beta$ -glucans in medicine, and the known mechanisms underlying their biological effects. We suggest why structurally different  $\beta$ -glucans may have different efficiencies of action, and how they might be better applied clinically, safely and effectively.

## Effects of $\beta$ -glucans on the immune system

Some fungal  $\beta$ -glucans markedly stimulating our immune system protect us from attack by pathogenic microbes and from harmful effects of environmental toxins and carcinogens (Vetvicka & Yvin 2004; Brown & Gordon 2005).  $\beta$ -Glucans are not synthesized by humans, so these compounds are recognized by our immune systems as non-self molecules, inducing both innate and adaptive immune responses (Fig 2) (Brown & Gordon 2005).

### Effects on the innate immune system

Innate immunity is present when we are born, and is a relatively non-specific system, responding to many, but not all, structurally related antigens (Brown & Gordon 2001; Munz et al. 2005). Certain  $\beta$ -glucans, including zymosan, grifolan (GRN) and lentinan (Table 1), appear to activate phagocytes, thus leading to elimination of pathogens by phagocytosis (Ladanyi et al. 1993; Kurashige et al. 1997; Brown et al. 2002). Among these, the macrophages preferentially attack dead cells and intracellular pathogens (Munz et al. 2005). Natural killer cells (NKs) circulate in blood to lyse cancer and virus-infected cells, whereas neutrophils are effective against pyogenic bacteria. Krestin (PSK), the protein-bound polysaccharide K containing  $\beta$ -(1 $\rightarrow$ 3)-glucans from *Coriolous versicolor* increases NK cytotoxic activity *in vitro* (Garcia-Lora et al. 2001; Garcia-Lora et al. 2003), whereas lentinan also activates type 3 complement receptors (CR3), which coat bacterial cells to facilitate their engulfment by phagocytes (Vetvicka et al. 1996; Vetvicka & Yvin 2004). In addition, zymosan (the cell wall particulate glucan from *Saccharomyces cerevisiae*), and MG [the microparticulate form of  $\beta$ -(1 $\rightarrow$ 3)-glucans from *S. cerevisiae*] both stimulate macrophages to produce cytokines, local immunomodulators, and





Fig 1 - Examples of structures of  $\beta$ -(1 $\rightarrow$ 3)(1 $\rightarrow$ 6) glucans showing branching patterns of their repeating units.

Table 1 - Examples of fungal  $\beta$ -glucans and their chemical diversity

Glucan	Abbreviation	Fungal Source	Structure	Reference
Curdlan		<i>Alcaligenes faecalis</i> (bacterial)	Linear $\beta$ -(1 $\rightarrow$ 3)-glucan	(McIntosh, Stone & Stanisich, 2005)
Grifolan	GRN	<i>Grifola frondosa</i>	Two $\beta$ -(1 $\rightarrow$ 6) branched glucose residues for every five $\beta$ -(1 $\rightarrow$ 3) glucose residues	(Munz, Steinman & Fujii, 2005)
Lentinan		<i>Lentinula edodes</i>	One $\beta$ -(1 $\rightarrow$ 6) branched glucose residue for every three $\beta$ -(1 $\rightarrow$ 3) glucose residues	(Sasaki & Takasuka, 1976)
Schizophyllan (Sizofiran)	SPG	<i>Schizophyllum commune</i>	One $\beta$ -(1 $\rightarrow$ 6) branched glucose residue for every three $\beta$ -(1 $\rightarrow$ 3) glucose residues	(Kikumoto S, 1971)
(Sonifilan)	SPG		Ultrasonic partial degradation of schizophyllan	(Johnson et al. 1963)
Scleroglucan	SSG	<i>Sclerotinia sclerotiorum</i>	One $\beta$ -(1 $\rightarrow$ 6) branched glucose residue for every three $\beta$ -(1 $\rightarrow$ 3) glucose residues	(Garcia-Lora et al., 2003)
Zymosan		<i>Saccharomyces cerevisiae</i>	Crude cell wall extract of genetically engineering yeast containing a mixture of $\beta$ -(1 $\rightarrow$ 3)(1 $\rightarrow$ 6) glucans and mannose	(Sato et al., 2003)
	MG	<i>Saccharomyces cerevisiae</i>	Microparticulate form of the $\beta$ -(1 $\rightarrow$ 3) glucan	
Betafectin	PGG	<i>Saccharomyces cerevisiae</i>	$\beta$ -(1 $\rightarrow$ 6) branched glucose residues of $\beta$ -(1 $\rightarrow$ 3) glucan	(Cheung & Modak, 2002)
Krestin	PSK	<i>Trametes versicolor</i>	Protein-bound $\beta$ -(1 $\rightarrow$ 3) glucan	(Kurashige, Akuzawa & Endo, 1997)
Yeast whole $\beta$ -glucan particules	WPG	<i>Saccharomyces cerevisiae</i>	Crude preparation $\beta$ -(1 $\rightarrow$ 6) branched glucose residues of $\beta$ -(1 $\rightarrow$ 3) glucans	(Lebron et al., 2003)
Pestolotan		<i>Pestalotia</i> sp. 815	Three $\beta$ -(1 $\rightarrow$ 6) branched glucose residues every five $\beta$ -(1 $\rightarrow$ 3) residues	(Misaki et al., 1984)
Epi-glucan		<i>Epicoccum nigrum</i>	Two $\beta$ -(1 $\rightarrow$ 6) branched glucose residues every three $\beta$ -(1 $\rightarrow$ 3) glucose residues	Schmid et al., 2001

Fig 2 - Immunostimulation by fungal  $\beta$ -glucans.

these in turn activate adaptive immunity (Fig 2), (Ladanyi *et al.* 1993; Kurashige *et al.* 1997) (Brown *et al.* 2002; Sato *et al.* 2003; Young *et al.* 2004). Zymosan can stimulate the production and activity of cytokines such as interleukin (IL)-2, IL-10 and IL-12 (Du *et al.* 2006; Brown 2006), and *in vitro* studies show that MG is taken up rapidly by peritoneal macrophages (Berner *et al.* 2005). Synthesis of mRNAs of cytokines, e.g. tumour necrosis factor alpha (TNF- $\alpha$ ), IL-6 and IL-1, is then up-regulated and their secretion increased (Berner *et al.* 2005). However, neither zymosan, PSK nor MG are chemically pure  $\beta$ -(1 $\rightarrow$ 3)-glucans, so it is possible that other chemical components in these preparations may be responsible. However, when chemically characterized zymosan, treated to produce the particulate form containing only  $\beta$ -(1 $\rightarrow$ 3)-glucan was added to cultured macrophage cells, the production of IL-10, reactive oxygen species (ROS) and TNF increased in a dose-dependent way (Saijo *et al.* 2007). Similarly, the addition of a pure glucan (SCG) from *Sparassis cripa* increased the production of IL-12, TNF and interferon gamma. Thus, increasing evidence is now available from studies with high purity  $\beta$ -glucans and a range of host cells that would seem to indicate that these are the effective immunomodulators.

### Effects on adaptive immunity

The adaptive immune system responds to introduced foreign antigens (Munz *et al.* 2005). It involves both B and T cells. The former produce antibodies to mediate humoral immunity, whereas T cells induce cell-mediated immunity (Munz *et al.* 2005). Cytokines promote T cell differentiation to helper T cells 1 (Th1) and 2 (Th2), which mediate cell and humoral immunities, respectively (Trinchieri 2003). The adaptive immune response also involves dendritic cells (DC), derived from monocytes, and these present antigens to T-cells for activation of immune responses (Munz *et al.* 2005). These DC are activated by PSK, thus facilitating the expression of adaptive immunity (Kanazawa *et al.* 2004; Munz *et al.* 2005).

Fungal  $\beta$ -glucan-induced immune responses are different in their actions to immune therapies based on supplementation of elements of the immune system, e.g. exposure to IL2 and interferon gamma. Instead, they appear to act by stimulating the whole immune system. Consequently, these  $\beta$ -glucans may have an advantage in treating diseases. Furthermore, many can be administered orally, and combinational application with other immune therapies may generate a more potent end result. For example, such a synergistic effect is believed to result from combining  $\beta$ -glucan application with either BCG or interferon gamma (Berner *et al.* 2005; Drandarska *et al.* 2005).

### Possible mechanisms of fungal $\beta$ -glucan action

Multicellular organisms possess receptors called 'pattern recognition receptors' (PRRS), to detect innately non-self structures (including pathogen-associated molecular patterns, or PAMPs) (Brown & Gordon 2005). Thus, fungal  $\beta$ -glucans probably act as PAMPs and are recognized by appropriate cell-surface receptors, initiating immune responses. In humans, a number of such receptors have been identified. These are dectin-1, complement receptor 3 (CR3), scavenger receptors,

lactosylceramide (LacCer), and the toll-like receptor (TLR). Their roles and consequences of their interactions with  $\beta$ -glucans are summarized in Table 2 and Fig 3 (Brown & Gordon 2005). Evidence suggests that dectin-1 is most important in the activation innate immune responses in macrophages (Herre *et al.* 2004a; Willment *et al.* 2005), as blocking with an anti-dectin-1 antibody and knockout of the dectin-1 gene resulted in the abolition of all macrophage-mediated responses (Steele *et al.* 2003; Taylor *et al.* 2007). Other receptors may also be involved. They include those detected in the U937 cell line, where two receptors other than CR3 could bind with scleroglucan, schizophyllan, laminarin, a glucan phosphate and a glucan sulphate (Mueller *et al.* 2000). However, their roles are much less clearly defined than that of dectin-1.

### Dectin-1

Dectin-1 is a lectin consisting of four components: an extracellular carbohydrate-recognition domain (CRD), a stalk, a trans-membrane region, and an intracellular cytoplasmic tail (Ariizumi *et al.* 2000; Brown & Gordon 2001; Marshall *et al.* 2004). Several human dectin-1 isoforms have been cloned and characterized (Ross 2000; Hermanz-Falcon *et al.* 2001; Willment *et al.* 2001; Yokota *et al.* 2001). Of these, BGRB and BGRB are the two major isoforms, with BGRB lacking the stalk (Willment *et al.* 2001). Dectin-1 is commonly expressed in macrophages, neutrophil lineages, DC, and some T-cells, but not in NK cells (Ross 2000; Herre *et al.* 2004a; Underhill *et al.* 2005; Willment *et al.* 2005).

The function of dectin-1 in binding to zymosan is achieved by the CRD, which also binds to intact fungal cells and other soluble fungal  $\beta$ -(1 $\rightarrow$ 3) and/or  $\beta$ -(1 $\rightarrow$ 6)-glucans (Ariizumi *et al.* 2000; Marshall *et al.* 2004). Dectin-1 consists of 244 amino acids, and has six cysteine residues, all of which are highly conserved (Ariizumi *et al.* 2000). Two amino acids (Trp221 and His223) located after the fourth cysteine residue appear to be critical in its binding function (Adachi *et al.* 2004). Zymosan binding may also require other compounds, like the CD63 tetraspanin receptor and pentraxin3, a tumour necrosis factor (Diniz *et al.* 2004; Mantegazza *et al.* 2004). Evidence has shown that dectin-1 binds specifically to  $\beta$ -(1 $\rightarrow$ 3)-glucans, but only those consisting of at least 10-mer oligosaccharides. However, the mechanism for this binding selectivity is not known (Palma *et al.* 2006). There is a need to identify more precisely the structural features of  $\beta$ -glucans that determine the subsequent level of activation of immune responses, and more studies with those whose chemical structures have been fully characterized are required.

Binding of dectin-1 with the ligand activates several signalling pathways to promote innate immune responses through activation of phagocytosis, ROS production, and induction of inflammatory cytokines (Willment *et al.* 2001; Grunebach *et al.* 2002). The cytoplasmic domain of dectin-1 has an immunoreceptor tyrosine-based activation motif (ITAM) to activate a tyrosine kinase, which in turn stimulates ROS production but not phagocytosis (Underhill *et al.* 2005; Rogers *et al.* 2005). Activation of this tyrosine kinase also induces synthesis of TNF- $\alpha$ , and IL-2, IL-10, IL-12.

Table 2 - Nature of the known receptors for  $\beta$ -glucans

Receptor type	Chemical nature	Affected cells	Effects on immune response	Other ligands
Dectin-1	Glycoprotein	Monocytes Macrophages Neutrophils DC cells T-cells	Phagocytosis Endocytosis ROS production Cytokines (TNF- $\alpha$ , IL-2, 10, 12) MIP-2,	Intact fungi, T-cells
CR3	A heterodimer Consist of CD11b and CD18 chains	Myeloid NK Neutrophils Lymphocytes	Tumour cytotoxicity	A variety of pathogens
Scavenger	A heterogeneous group of molecules	Myeloid Endothelial	Immunity	LDL HDL Polyanionic Microbes
LacCer	Glycosphingolipid containing a hydrophobic ceramide lipid and a hydrophilic sugar	Neutrophils Epithelial	ROS Anti-microbial Cytokines (NF- $\kappa$ B) Lyn kinase	A variety of microbes
TLR	A novel protein family with associated protein MyD88	Macrophages DC Lymphocytes Epithelial	Cytokines (NF- $\kappa$ B, TNF- $\alpha$ , IL12)	A wide spectrum of microorganisms


CR3, complement receptor 3; CRD, carbohydrate-recognition domain; DC, dendritic cell; eNOS, endothelial nitric oxide synthase; LacCer, lactosylceramide; IL, interleukine; MIP, macrophage inflammatory protein-2; NK, natural killer cell; NF- $\kappa$ B, nuclear factor-kappaB; ROS, reactive oxygen species; TNF- $\alpha$ , tumour necrosis factor-alpha.

Several pathways have now been identified as being involved in dectin-1 downstream signalling (Fig 3). First, some evidence suggests it might act synergistically with TLR to produce strong inflammatory responses by stimulating cytokines such as TNF- $\alpha$ , IL-2 and IL-12 (Gantner et al. 2003; Brown 2006). Although both dectin-1 and TLRs are activated by zymosan (Rogers et al. 2005), the TLR ligand is not yet known. The suggestion is that it may not be a  $\beta$ -glucan as laminarin had no effect on its signalling pathway, and hot alkali treated zymosan only bound to dectin-1 (Gantner et al. 2003). Recent evidence also showed that SCG and NaClO-oxidized zymosan (which contained only  $\beta$ -glucan) induced production of TNF and IL-12 and interferon gamma was not affected by a deficiency of MyD88, indicating its effect is independent of the TLR pathway (Saijo et al. 2007). Whether other fungal  $\beta$ -glucans can bind to TLR is not known. This needs to be determined, and it could be that this synergistic effect arises because phosphorylated dectin-1 can form a complex with TLR (Brown 2006).

In addition, another pathway independent of TLR is mediated via spleen tyrosine kinase (Syk) to produce other cytokines, including the macrophage inflammatory protein-2 (MIP2, CXCL2) and IL-2 and IL-10 in mice DC cells (Rogers et al. 2005). After binding to the ligand, dectin-1 is phosphorylated by a non-receptor tyrosine kinase Src. Syk is then activated, which in turn activates the card9-bcl10-Malt1 complex (caspase recruitment domain 9 - B-cell chronic lymphocytic leukaemia/lymphoma 10-mucosa-associated lymphoid tissue lymphoma translocation gene 1). This complex mediates the induction of cytokines like nuclear factor (NF)- $\kappa$ B and IL production. Zymosan failed to activate NF- $\kappa$ B, TNF- $\alpha$ , IL-2 and

IL-6 production in card9-/- and Malt1-/- mouse dendritic cells indicating the importance of the card9-bcl10-Malt1 complex in this pathway (Gross et al. 2006). Consequently, animals with Card9-/- were much more susceptible to *Candida albicans* infections (Gross et al. 2006). Surprisingly, dectin-1-deficient mice had a similar capability to counteract *C. albicans* infections, indicating that another pathway to activate card9 by  $\beta$ -glucan must exist (Saijo et al. 2007). There is now evidence that CR3 also activates Syk (Li et al. 2006).

Furthermore, phagocytosis in macrophages is another signalling pathway activated by the dectin-1 receptor (Underhill

Fig 3 - Possible fungal  $\beta$ -glucan mediated signal pathways.

et al. 2005; Brown 2006) which seems to be independent of any involvement by either TLR or Syk. The dectin-1 cytoplasmic domain has three consecutive acidic amino acids signalling phagocytosis and activation of phagocyte lysosomal endosomes (Engering et al. 2002; Serrano-Gomez et al. 2004; de la Rosa et al. 2005). However, its detailed mode of action is less clear. Dectin-1 has also been shown to mediate zymosan-induced arachidonic acid production and cyclooxygenase 2 expression through its cytoplasmic domain. Both promote acute tissue inflammation (Suram et al. 2006).

### CR3 receptor

The CR3 receptor, consisting of CD11b and CD18 domains, recognizes a wide range of microbial cells, and functions as an adhesion molecule. It is expressed mainly on neutrophils, monocytes and NK cells, but not macrophages (Ross 2000). Two binding sites exist in CD11b. One is for  $\beta$ -glucans, and is located within the C terminus, while the other for iC3b (cleaved component 3 fragment of serum complement system), is located within the N-terminus (Thornton et al. 1996; Xia & Ross 1999).

Binding of  $\beta$ -glucans to CR3 increases adhesion to microbial cells and activates the iC3b pathway causing tumour cytotoxicity (Xia & Ross 1999; Xia et al. 2002) (Tsikitis et al. 2004; Kanse et al. 2004). Such activation requires occupation of both binding sites as cytotoxicity is blocked by an anti-CR3 antibody (Vetvicka et al. 1996; Li et al. 2006). Zymosan increased NK cell-mediated killing of tumour cells via activation of CR3 (Vetvicka & Yvin 2004; Diniz et al. 2004). However, the trials on which this conclusion was based were only conducted in neutrophils and NK cells, and other cells may possess different receptors. NK cells have no dectin-1 receptors, so CR3 may be the major receptor. Neutrophil cells have both CR3 and dectin-1 receptors, but is it not known what happens when CR3 receptors are blocked, or what the specific role of dectin-1 is. Zymosan also binds to dectin-1 in neutrophil cells to activate immune responses (Willment et al. 2005), and if an anti-CR3 antibody inhibits these, then it is possible that dectin-1 and CR3 are both needed for zymosan-induced activation.

### Scavenger receptors

Scavenger receptors located in myeloid and endothelial cells comprise a heterogeneous group of proteins with two transmembrane domains, two intracellular domains and one extracellular domain (Acton et al. 1994; Rice et al. 2002; Assanasen et al. 2005). These recognize a range of foreign cells, low-density lipoprotein (LDL), high-density lipoprotein (HDL) and selected polyanionic ligands (Assanasen et al. 2005). Although lentinan can bind to these receptors, no specific  $\beta$ -glucan scavenger receptors have yet been identified (Rice et al. 2002). Multiple signalling pathways appear to be activated by the Src receptor, including those involving Src family kinase(s), phosphatidylinositol-3 kinase (PI3K), Akt kinase, and p38 mitogen-activated protein kinase (MAPK), and an endothelial nitric oxide synthase (eNOS) (Mineo et al. 2003; Assanasen et al. 2005). However, their possible role, if any, in

the biological effects mediated by fungal  $\beta$ -glucans is still unknown, and no evidence is available that they are important.

### LacCer

LacCer located in neutrophil and endothelial cells is a glycolipid containing a hydrophobic ceramide lipid and a hydrophilic sugar moiety. It recognizes both microbial cells and fungal  $\beta$ -(1 $\rightarrow$ 3)-glucans (Zimmerman et al. 1998). In alveolar epithelial cells, the  $\beta$ -(1 $\rightarrow$ 3)-glucan cell wall component of *Pneumocystis carinii* increased synthesis of NF- $\kappa$ B, MIP-2 and TNF- $\alpha$  through a protein kinase C signalling pathways (Wang et al. 2005; Evans et al. 2005), while in neutrophil cells, LacCer increased ROS production to kill microbes through activation of PI-3K, MAPK, and protein kinase C mediated by the Src family kinase Lyn (Iwabuchi & Nagaoka 2002). The role played by  $\beta$ -(1 $\rightarrow$ 3)-glucans in mediating these immune responses again requires further study, preferably in experiments using purified preparations of known chemical structure.

### TLRs

TLRs are transmembrane receptors of a novel protein family. At least 11 members of this family exist in humans (Roeder et al. 2004). They respond to the presence of a diverse group of microbes including fungi, bacteria, viruses and protozoa (Roeder et al. 2004). Fungal cells and zymosan bind to TLR 2 and 4, and both activate innate immune responses (Takeda et al. 2003). TLR2 causes an increase in the levels of NF- $\kappa$ B and cytokine production, including TNF- $\alpha$  and IL-12, which is mediated by the adaptor protein MyD88 (Ozinsky et al. 2000; Kataoka et al. 2002; Lebron et al. 2003). However, most of these data were again generated in studies with chemically crude  $\beta$ -glucans. For example, the zymosan used probably contained the lipopolysaccharide (LPS) endotoxin from Gram-negative bacteria, which is also a strong inducer of TLRs (Kawai et al. 2001; Duenas et al. 2006; Lapaque et al. 2006).

In summary, what seems clear is that recognition of  $\beta$ -glucans by these receptors leads to elimination of bacterial pathogens, and a single soluble  $\beta$ -glucan preparation may activate several receptors, as seen with zymosan which binds to dectin-1 and CR3 (Ariizumi et al. 2000; Diniz et al. 2004; Takeda et al. 2003). Fungal  $\beta$ -glucans may activate different immune cell populations through specific receptors, although sometimes several receptors may exist in a single cell. Dectin-1 seems to be most important in macrophages, and while CR3 has a role in NK cells, different receptors are present in neutrophils. Although co-ordinated effects between dectin-1 and TLR are well documented, those involving other receptors have not been well studied.

### Importance of $\beta$ -glucans in the treatment of cancer

Since their anti-tumour activity was first demonstrated nearly 50 y ago, many animal experiments have demonstrated the remarkable effects of certain fungal  $\beta$ -glucans on a range of tumours (Vetvicka & Yvin 2004). Several human clinical trials have also shown possible treatment benefits, although these data are still preliminary and controversial. Currently,

lentinan, schizophyllan, and PSK are approved in Japan for clinical use in human cancer treatment (Mizuno et al. 1999).

Cancer is uncontrolled cell proliferation induced by many factors including environmental chemicals, viruses, bacteria, hormones, and chronic inflammation (Nam et al. 2005). Three developmental stages are recognized (Borchers et al. 2004; Nam et al. 2005). The first is initiation, in which a mutagen binds to the cell DNA and causes damage, which by itself is usually insufficient to induce tumour production. The second stage is activation of a tumour promoter that leads to the formation of small benign tumours. Finally in the third stage, progression, the normal tight control over the cell cycle is lost, resulting in uncontrolled cell proliferation (Borchers et al. 2004). Studies with oligonucleotide microarrays in hepatocellular carcinoma have revealed that multiple gene mutations are involved in tumour development (Nam et al. 2005). Thus, as many as 3084 altered genes correlated with tumour progression from the formation of dysplastic nodules eventually to overt hepatocellular carcinoma. Among these, 240 genes were associated with different stages in tumour development.

Although surgical resection remains the most effective early treatment of solid cancers, chemotherapies and immunotherapies are often used (Nilsson et al. 2004). Progress in understanding the molecular mechanisms of carcinogenesis has allowed these non-surgical therapies to become more effective. Some fungal  $\beta$ -glucans appear to beneficially influence both cancer promotion and progression (Takaku et al. 2001; Nilsson et al. 2004; Nam et al. 2005), and such treatment of rats has led to formation of much smaller tumours than those seen in controls (Vetvicka & Yvin 2004).

Fungal  $\beta$ -glucans also have synergistic effects with monoclonal antibodies used in cancer treatment. Monoclonal antibody therapy targets key components of the biological pathways involved in carcinogenesis. Furthermore, yeast  $\beta$ -glucans given orally with monoclonal antibody therapy increased neuroblastoma tumour regression and long-term survival in mice (Cheung et al. 2002; Cheung & Modak 2002; Yan et al. 2005). In mice with established subcutaneous non-Hodgkin's lymphoma xenografts, a combination of intravenous complement-activating antibody, rituximab, and WGP from yeast had a higher therapeutic efficacy than treatment with either alone (Modak et al. 2005). WGP induced the killing ability of Mab G250, an antibody against the enzyme carbonic anhydrase, and present only in renal carcinoma cells, (Sier et al. 2004), by assisting granulocytes to kill iC3b-coated cancer cells. Intravenously and orally administered  $\beta$ -glucans (e.g. WPG) promoted tumour regression by stimulating granulocytes and macrophages, as well as triggering cytotoxicity of tumour cells, but not in CR3<sup>-/-</sup> mice, indicating a critical role for CR3 (Allendorf et al. 2005; Yan et al. 2005). Two domains of the CR3 receptor must both be occupied by iC3b and the  $\beta$ -glucan to exert this killing effect (Sier et al. 2004; Li et al. 2006). As no  $\beta$ -glucans are synthesized by humans, monoclonal antibodies enhance treatment capabilities only after their administration (Sier et al. 2004).

Although these data are impressive, they are not easy to interpret mechanistically, as none of the  $\beta$ -glucans used in the studies were confirmed as being chemically pure. In addition, some chemical variation in the structure of most natural

products is to be expected. Explanations for their mechanism of action are also limited to considering a role for the CR3 receptor alone.

Thus, concomitant use of the anticancer drug 5-1 and lentinan prolonged survival times of colon-26-bearing mice (Mushiake et al. 2005), but not in athymic mice, which suggests some role for DCs. These mice showed increased CD86<sup>+</sup> DCs infiltration into colon-26 and more potent activities for T-cell proliferation and T-lymphocyte cytotoxicity. Consequently, DC may play a key role in stimulating anti-tumour immunity in combination with lentinan. Orally administered WGP may be taken up by macrophages in the gastrointestinal tract and shuttled to the bone marrow where it is degraded and secreted as smaller soluble  $\beta$ -(1 $\rightarrow$ 3)-glucan molecules, which then bind to the receptor CR3 (Hong et al. 2004). Further consideration will be given to the possible metabolic fate of ingested  $\beta$ -glucans later.

Certain fungal  $\beta$ -glucans may also ameliorate chemotherapy and radiation treatment by increasing patient tolerance, and speed recovery from toxic effects (Harada et al. 2002; Gu et al. 2005), the most severe of which is leukopenia, associated with an increased risk of infection. In a mouse model with cyclophosphamide-induced leukopenia, oral administration of the  $\beta$ -glucans from *Sparassis crispa* markedly increased the rate of leukocyte recovery (Harada et al. 2002). Furthermore, intraperitoneal injection of a yeast derived  $\beta$ -glucan decreased mortality in tumour-bearing mice exposed to whole-body X-ray radiation, and increased their leukocyte and lymphocyte numbers, including both NK and lymphokine-activated killer (LAK) cells (Gu et al. 2005).

PSK has been successful in postoperative treatment of resectable cancer in humans, increasing survival rates (Munemoto et al. 2002; Ito et al. 2004). A literature search has revealed 48 published clinical trials using PSK (Sakamoto et al. 2006). Three involved 1094 patients, and results showed that adjuvant immunochemotherapy with PSK increased five-year overall survival rates from 72.2% to 79%, and five-year disease-free rates from 65.9% to 72.2% (Nakazato et al. 1994; Ito et al. 2004; Ohwada et al. 2004). Although both sets of data are statistically significant, in the trial of Ito et al. no difference was recorded for seven-year overall survival and disease-free rates (Ito et al. 2004). Thus, these data are still very preliminary; again the major weakness in interpreting these data is that PSK is not a chemically pure  $\beta$ -glucan, but a protein (25%)-bound polysaccharide extracted from mycelium of *C. versicolor* (Kobayashi et al. 1995). It is thought to consist of a main chain of  $\beta$ -((1 $\rightarrow$ 4)-linked glucose residues with a side-chain of  $\beta$ -(1 $\rightarrow$ 3) as well as  $\beta$ -(1 $\rightarrow$ 6) linked glucose residues. There is also the possibility that PSK, like WSP, may be degraded to smaller molecules in the body, which then bind to receptors.

Lentinan has also been used in similar clinical trials and been shown to prolong survival times of humans with gastric cancer from 199 to 297 d of the median survival period (Nakano et al. 1999). A crude  $\beta$ -(1 $\rightarrow$ 6)-glucan extracted from *Agaricus blazei* basidiocarps caused apoptosis or programmed cell death in human ovarian cancer HRA cells. As the apoptotic effect of this  $\beta$ -glucan could be abolished by applying the p38 MAPK-specific inhibitor, SB203580 (Kobayashi et al. 2005), this would seem to involve activation of the p38 MAPK

pathway by translocation of an apoptosis activator Bax from the cytosol to the mitochondria, cytochrome c release and caspase 9 activation. In our opinion, these clinical trials are still very preliminary, and the data from them should be treated cautiously.

In addition, lentinan, GRN, SSG, and SPG also reduce preventatively the overall risks of cancer (Greenwald et al. 2001). As these are caused by mutations, oxidative stress, and inflammation, and develop in several stages from multiple gene mutations, they are preventable by blocking the appropriate regulatory molecules involved in these processes. These are known to include Nrf2 (nuclear factor erythroid 2-related factor 2), epidermal growth factor receptor kinases, phosphatidylinositol 3-kinase, components of the Janus kinase-signal transducers and activators of transcription (JAK-STAT) pathway, NF- $\kappa$ B and cyclin D (Sporn & Liby 2005). Cancers examined include melanoma, adenocarcinoma, mammary carcinoma, lymphocytic leukaemia, and Lewis lung carcinoma (Suzuki et al. 2001). The most impressive data come with lentinan, which in animal models prevented carcinogenesis in chronic ulcerative colitis by inhibiting expression of P450 1A2, which catalyses pre-carcinogenic compounds (Mitamura et al. 2000; Okamoto et al. 2004). This inhibition was mediated by an increase in TNF- $\alpha$  levels and DNA-binding activity of the NF- $\kappa$ B. Other fungal  $\beta$ -glucans of known chemical composition should be screened to see whether common mechanisms of action might exist.

In summary, fungal  $\beta$ -glucans may exert multiple effects on cancer cells and prevent cancer. Their mechanisms of action appear to be complex. The regulation by  $\beta$ -glucans of the signalling pathways of tumour production involves many key elements (Sporn & Liby 2005). Most studies so far demonstrate that  $\beta$ -glucans cause regression in tumour size, but not total tumour eradication. These are different because regressed tumours can enlarge rapidly when treatment stops.

### Importance of fungal $\beta$ -glucans in the treatment of infectious diseases

The major current problem in controlling infectious diseases is antibiotic resistance (Suzuki et al. 2001). Some fungal  $\beta$ -glucans are effective against almost all known pathogenic microbes, by acting through several different immunomodulatory mechanisms.

#### Anti-viral activity of fungal $\beta$ -glucans

SPG, lentinan, zymosan, and bacterial curdlan all appear effective against several viral agents (Mayell 2001), although most data available are only descriptive. SPG controls chronic hepatitis B infections, by modulating both cellular and humoral immune responses (Kakumu et al. 1991). In mice injected with hepatitis virus strain MHV-A 59, a  $\beta$ -glucan prepared from *Saccharomyces cerevisiae* cells increased infected cell survival by inhibiting cell necrosis and activating phagocytic activity (Vetvicka & Yvin 2004).

Zymosan from *S. cerevisiae* enhanced the immune response in humans administered the human immunodeficiency virus (HIV) vaccine by stimulating Th cell-mediated

immunity through activation of the complement system and interferon gamma (Ara et al. 2001). The same  $\beta$ -glucan also decreased viral nucleic acid levels in cells of pigs infected with swine influenza virus, and led to increases in their interferon gamma and nitric oxide levels (Jung et al. 2004).

Some preliminary clinical trials have been conducted with lentinan in the treatment of HIV patients suffering no accompanying opportunistic infections. In a placebo-controlled trial in San Francisco, patients (ten per group) with HIV were administered 2, 5, and 10 mg lentinan or the placebo intravenously. Results showed increased CD4 cell numbers in those administered lentinan (Gordon et al. 1998). When lentinan was applied with didanosine in HIV-positive patients with CD4 levels of 200–500 cells mm<sup>-3</sup>, CD4 cell numbers increased by about 142 CD4 cells mm<sup>-3</sup> over 12 m, whereas didanosine alone led to decreased CD4 numbers (Gordon et al. 1995). These trials suggest that lentinan was more effective in combination with other antiviral agents, although the patient numbers used were small.

#### Anti-bacterial activities of fungal glucans

Some  $\beta$ -glucans, including lentinan, WGP, PPG, and SSG, are also effective against bacterial infections. Thus, lentinan reduced *Mycobacterium tuberculosis* infections by increasing macrophage levels *in vivo* in a rat model and *in vitro* examinations showed these macrophages had increased killing ability toward *M. tuberculosis* cells (Markova et al. 2003; Markova et al. 2005). Both PPG and WGP have effectively treated mice against *Bacillus anthrax* infections (Kournikakis et al. 2003), correlating with increases in levels of cytokines IL-2 and IL-10, although experiments to clarify their modes of action by blocking cytokines were not conducted. When SSG was tested against *Streptococcus pneumoniae* types 4 and 6B in mice infected experimentally intraperitoneally (Hetland et al. 2000), it protected animals against both strains when applied 3 d before challenge. However, it was only effective against the type 4 strain when administered after infection.

PGG from *Saccharomyces cerevisiae* also effectively treated *Staphylococcus aureus* infections involving strains resistant to several  $\beta$ -lactam antibiotics including methicillin, by improving patient survival by 80% (Liang et al. 1998). It also successfully treated methicillin-resistant strains of *S. aureus* and *S. epidermidis* in guinea pigs (Kernodle et al. 1998). As with other examples given here, only a restricted range of  $\beta$ -glucans have been examined in this way and further work with others (Table 1) seems warranted. PGG can also prevent wound infections, and was more effective in combination with the antibiotic cefazolin than alone (Kaiser & Kernodle 1998). However, no explanation was provided.

When SPG and zymosan were examined for their effects on tissue injury induced by LPS, the outcomes were controversial. The claim that SPG reduced LPS-induced toxicity both in rats and perfused liver (Kukan et al. 2004) by reducing the basis for LPS toxicity that requires induction of TNF- $\alpha$  at a transcriptional level (Harrison et al. 2004), has been questioned. Zymosan may also act by stimulating hepatic macrophages (Kupffer cells) and increasing their sensitivity to LPS, leading to increased TNF- $\alpha$ -induced hepatic lesion (Beutler & Cerami 1989; Yamaguchi et al. 2001).

Overall, the evidence seems to suggest that  $\beta$ -glucan therapy should be better based on disease prevention than its cure, where the immune system can be stimulated first to cope better with subsequent infections. No studies have been performed yet to examine the possible effect of dectin-1 and other receptors in this antibacterial role of fungal  $\beta$ -glucans, although it seems unlikely that different receptors would be involved.

#### Anti-fungal activity of fungal $\beta$ -glucans

Currently, opportunistic fungal infections are problematic, especially in immunocompromised patients with HIV and cancer, so any stimulation of the immune system may be critical in patient survival. As mentioned earlier,  $\beta$ -glucans are major components of walls of pathogenic fungi and probably act as important inducers of immune responses against them by binding to the dectin-1 receptor (Netea et al. 2006; Saijo et al. 2007; Taylor et al. 2007). Some edible  $\beta$ -glucans can activate phagocytic cells including macrophages, neutrophils and DC to enhance the host's innate response to fungal infections (Herre et al. 2004b). For example, MG up-regulated synthesis of the cytokines TNF- $\alpha$ , IL-6 and IL-1 mRNA in macrophages, neutrophils, and DCs (Bernier et al. 2005). Several  $\beta$ -glucan receptors appear to be involved. Cell wall  $\beta$ -glucans of *Pseudomonas carinii* bind to dectin-1 and Lac receptors on alveolar epithelial cells to promote innate immune responses (Hahn et al. 2003). The signalling protein MIP2 is activated and cytokines then released. A synergy exists between interferon-gamma and these wall  $\beta$ -glucans, as pre-treatment with interferon-gamma increases the sensitivity of response. Other  $\beta$ -glucans like zymosan can protect host cells against *Candida albicans* and *P. carinii* infections, involving their recognition by dectin-1 (Hobson et al. 2004; Gantner et al. 2005; Steele et al. 2003; Viriyakosol et al. 2005). TLR has also been shown to be involved in host cell immune responses to some fungal infections (Roeder et al. 2004).

From *in vitro* experiments, peritoneal macrophages from mice treated with the particulate  $\beta$ -glucan from *Saccharomyces cerevisiae* showed enhanced activity against *P. brasiliensis* infections (Pelizon et al. 2005). Their spleen cells had higher synthetic abilities for TNF- $\alpha$  and IL-12, and IL-12 treatment can inhibit spread of *P. brasiliensis* cells to the liver and spleen (Calich et al. 1998; Arruda et al. 2002). In a monocyte monolayer cell culture, both mRNA and cytokine TNF- $\alpha$  and IL-1  $\beta$  were induced by purified yeast glucan particles (Abel & Czop 1992), an event that was partially preventable by trypsin exposure, indicating a possible involvement of trypsin-sensitive  $\beta$ -glucan receptors, which were not identified.

Treatment of opportunistic fungal infection by traditional chemotherapy is not very effective. However, combinational therapy with antifungal agents and the immune stimulator granulocyte colony-stimulating factor (G-CSF) has successfully treated immunocompromised mice infected with *Aspergillus fumigatus* (Sionov et al. 2005). Fungal  $\beta$ -glucans have not yet been incorporated into such treatment regimes, although theoretically there seems no reason why they should not act synergistically with other anti-fungal agents and immune stimulators.

## Role of fungal $\beta$ -glucans in the treatment of hypercholesterolaemia, diabetes, high blood pressure, and wound healing

### Hypercholesterolaemia

Cardiovascular disease related to elevated blood cholesterol levels is still the most common cause of death in humans in western countries (Strong et al. 2005). Cholesterol contains LDLs and HDLs (Strong et al. 2005). LDL has been associated with an increased risk of coronary artery disease. Yeast  $\beta$ -glucans appear to be effective in lowering blood cholesterol concentrations, but the mechanism by which this occurs is still unclear (Nicolosi et al. 1999); there are few studies that have tested other fungal  $\beta$ -glucans for this activity.

### Diabetes

Several fungal  $\beta$ -glucans may reduce blood glucose concentrations after eating, possibly by delaying stomach emptying so that dietary glucose is absorbed more gradually (Lo et al. 2006). Orally ingested fruiting bodies and the acidic polysaccharide of both *Tremella mesenterica* and *T. aurantia* both reduced blood glucose concentrations in induced diabetic rats (Kiho et al. 1995). However, the structure of the active polysaccharide component was later shown to have  $\alpha$ -(1 $\rightarrow$ 3)-linked D-mannopyranosyl residues as the backbone and (1 $\rightarrow$ 3)-linked  $\beta$ -D-xylopyranose side chains at position 2, and  $\beta$ -D-glucopyransyluronic acid at position 4 (Kiho et al. 2000), so it was not a  $\beta$ -glucan. This emphasizes the risks associated with interpreting data from experiments where crude fungal preparations are used. A crude exopolysaccharide produced from submerged mycelial cultures of *Phellinus baummi* also exhibited hypoglycaemic effects in streptozotocin-induced diabetic rats (Kiho et al. 2000), but again the active component was not convincingly verified as a  $\beta$ -glucan. In genetically diabetic mice fed 20% whole mushroom maitake powder and its chemically derived fractions prevented an increase in blood glucose levels by increasing insulin sensitivity (Mayell 2001). A  $\beta$ -glucan prepared by hot water extraction of *Agaricus blazei* basidiocarps showed anti-hyperglycaemic, anti-hypertriglyceridaemic, anti-hypercholesterolaemic and anti-arteriosclerotic activity in diabetic rats, although the active component was not identified (Kim et al. 2005). When these preparations were digested by an endo  $\beta$ -(1 $\rightarrow$ 6)-glucanase from *Bacillus megaterium*, the resulting di- and tri-saccharides had double the anti-diabetic activities shown by the parent  $\beta$ -glucans, again raising question as to whether  $\beta$ -glucans or their derived oligosaccharides are the effective agents. No purified fungal  $\beta$ -glucan whose chemical structure has been fully characterized has yet been tested against diabetes. Consequently, the need is to clarify the structural features required and to identify the nature of the  $\beta$ -glucan binding receptors related to such activities if its promise is to be realized.

### Blood pressure

Some fungal  $\beta$ -glucans may also control blood pressure. An intravenous infusion of an aqueous extract of *Pleurotus sajor-caju*

mycelium caused a decrease in the mean systemic blood pressure in rats, although the biologically active component was not identified (Tam et al. 1986). However, in genetically deficient rats with spontaneous hypertension (SHR), a diet containing 5% maitake from *Grifola frondosa* had the same effect (Kabir et al. 1987; Kabir & Kimura 1989). Because of the complex chemistry of these basidiocarps, attempts have been made to elucidate which chemical components are responsible, and data again suggest that these may not always be  $\beta$ -glucans. Thus, in one trial a lectin from *Tricholoma mongolicum* basidiocarps appeared to be the effective agent (Wang et al. 1996). Responses with whole maitake basidiomes have also been compared with those from their ether-soluble and water-soluble extracts (Talpur et al. 2002), and as only consumption of the ether-soluble extract decreased systolic blood pressure in SHR, any role for  $\beta$ -glucans seems improbable. Furthermore, the component from Tamogi-take mushroom (*Pleurotus cornucopiae*) that inhibited angiotensin I and so decreased blood pressure in SHR was shown to be D-mannitol, (Hagiwara et al. 2005). Yet consumption of whole maitake basidiocarps and the water-soluble extract also led to a decrease in blood pressure in Zucker fatty rats, a diabetes rat model (Talpur et al. 2002). Thus, the possible involvement of fungal  $\beta$ -glucans in reducing hypertension is not yet clear, and as with many of the biological effects discussed here, now seems the time to direct our efforts to assessing more carefully purified, and structurally characterized individual fungal  $\beta$ -glucans.

#### Wound healing

Macrophage activities stimulated by  $\beta$ -glucans from *Saccharomyces cerevisiae* may also benefit wound healing and reduce scar tissue levels after surgery or trauma, as revealed by both animal and human studies (Mayell 2001; Portera et al. 1997). In normal human dermal fibroblasts this  $\beta$ -glucan preparation stimulated procollagen mRNA and collagen biosynthesis, together with increased NF-1 (Wei et al. 2002b). Inhibition of NF-1 by pentifylline blocked induction of procollagen mRNA by the same  $\beta$ -glucan, which also induced mRNA synthesis of many other wound growth factors including activator protein-1, specificity protein-1, neurotrophin 3, platelet-derived growth factor A, B, fibroblast growth factor acidic, fibroblast growth factor basic, transforming growth factor  $\alpha$ ,  $\beta$ , and vascular endothelial growth factor (Wei et al. 2002a). However, how the synthesis of these factors is regulated in the signal transduction pathway is unclear.

#### Immunomodulation by fungal $\beta$ -glucans: structure-function relationships

Traditionally the beneficial effects described here were reported in humans eating fungal basidiocarps, and not from consumption of single pure  $\beta$ -glucans, so most of the available information on relationships between structure and function is confused by this. However, it seems clear that individual fungal  $\beta$ -glucans differ in their effectiveness as immunomodulators. Even  $\beta$ -glucans with similar reported structures, molecular weights and solution conformations

can differ markedly. This may reflect our inability to acquire sufficient detail from their structural analyses to allow us to recognize possible subtle structural differences between them, using the present methodologies of nuclear magnetic resonance (NMR) and methylation analyses to determine branching patterns. Yet these response differences are much more noticeable when structurally quite different glucans are compared. The most popular  $\beta$ -glucans, lentinan, schizophyllan, grifolan, and SSG from *Lentinus edodes*, *Schizophyllum commune*, *Grifola frondosa*, and *Sclerotinia sclerotiorum*, are all  $\beta$ -(1 $\rightarrow$ 3)(1 $\rightarrow$ 6)-glucans, but with different reported branching frequencies (Table 1). All are effective against the same tumour model, but at quite different doses. Some papers have reported correlations between  $\beta$ -glucan effectiveness and molecular structure, size, branching frequency, structural modification, conformation and their solubility, but it is still difficult to confidently make generalizations because of the often contradictory data available. Furthermore, as mentioned earlier in this review we do not understand well the metabolic fate of ingested  $\beta$ -glucans, and whether they require modification and possibly conversion to smaller oligosaccharides, to have an effect. These are difficult questions to address experimentally, but the acidity of the stomach may readily generate partial acid hydrolysis products from them.

#### Molecular structure and size

The anti-tumour effects of fungal  $\beta$ -glucans seem to be related to their structural features. For example, a  $\beta$ -(1 $\rightarrow$ 3)-linked backbone seems essential (Seviour et al. 1992; Demleitner et al. 1992). So, although schizophyllan showed anti-tumour activity against Sarcoma 180 tumour cells, all glucans tested with  $\alpha$ -(1 $\rightarrow$ 3)-glucosidic links lacked any biological activity. Misaki et al. (1981) also showed that having  $\beta$ -(1 $\rightarrow$ 6)-linked side chains of glucose residues increased anti-tumour activity. Only a relatively small number of glucans have been examined, but most subsequent work has focused mainly on  $\beta$ -(1 $\rightarrow$ 3)(1 $\rightarrow$ 6)-glucans. However, factors other than their branching frequency may be important. Of these, molecular size seems to affect biological activity. Thus, for scleroglucan, only preparations of  $<50 \times 10^4 \text{ g mol}^{-1}$  or  $>110 \times 10^4 \text{ g mol}^{-1}$  were effective (Falch et al. 2000). Conversely, only low molecular weight lentinan had high anti-tumour activity (Zhang et al. 2005), so generalizations of this kind are risky.

#### Solubility of $\beta$ -glucans

The solubility of  $\beta$ -glucans depend on their degree of polymerization and thus their physical organization (Zekovic et al. 2005). Unsurprisingly, soluble  $\beta$ -glucans appear to be stronger immunostimulators than insoluble ones (Di Luzio et al. 1979; Di Luzio et al. 1980; Xiao et al. 2004), although the reasons are not totally clear, and orally administered insoluble  $\beta$ -glucans may be subsequently degraded into smaller bioactive oligomers after ingestion (Hong et al. 2004).

#### Branching frequency

The branching frequencies of fungal  $\beta$ -(1 $\rightarrow$ 3)(1 $\rightarrow$ 6)-glucans are thought to determine their biological activity (Seviour

et al. 1992), and a branching frequency of 0.2 (1 in 5 backbone residues) to 0.33 (1 in 3 backbone residues) was suggested as being optimal (Miyazaki et al. 1979). Thus, although both unbranched bacterial curdlan and several  $\beta$ -linked linear synthetic carbon 4 and 5 oligosaccharides (Jamois et al. 2005) showed some biological activities, chemical addition of  $\beta$ -(1 $\rightarrow$ 6)-linked glucose residues to the curdlan backbone led to increases in its activity against mice sarcoma 180 cells (Kihō et al. 1998). Binding studies with receptors in U937 cells suggested that the more highly branched scleroglucan, and schizophyllan had a higher affinity than that shown by the infrequently branched laminarin from the alga *Laminara digitata* (one every ten), but this in turn had a higher affinity than either linear glucan phosphate or glucan sulphate preparations (Mueller et al. 2000). In addition, scleroglucan has a much higher receptor binding ability than schizophyllan, although they reportedly have the same branching frequencies (Mueller et al. 2000). In this study the chemical structures of both  $\beta$ -glucan preparations were meticulously confirmed. Yet the important question of how and why branching frequencies might influence the host immune responses is not clear. It is also possible that the reported branching frequencies of their repeating units represent only the molecular average and that a considerable variation in it might occur within individual long  $\beta$ -glucans chains.

#### Branch modification

Some published information is available on how modifying the glucan side chains might affect their biological activity, and in some, it may increase it. When insoluble scleroglucan and its sulphated, carboxymethylated, methylated, hydroxyethylated and hydroxypropylated derivatives were tested against sarcoma 180 tumour and gastric carcinoma cells (Misaki et al. 1984; Wang et al. 2004), the sulphated and carboxymethylated forms showed the highest activity. The parent form had no anti-tumour activity against these cancers. Some evidence suggests that increased anti-tumour activity can also be achieved by polyol modification of the side chains residues of pestalotan (Misaki et al. 1984); but until more similar studies are performed with a greater diversity of glucans, any potential benefits from this approach will remain unclear.

#### Helical conformation

$\beta$ -Glucans can adopt either a single or triple helical conformation in solution, where hydrogen bonds hold the individual polymer chains together. Those with a triple helical configuration have been regarded as being the more powerful immunomodulators (Falch et al. 2000). For example, in a study by Falch et al. scleroglucan was only biologically active in a linear triple helical arrangement, and its subsequent denaturation reduced its cytokine inducing activity in monocytes (Falch et al. 2000). However, with schizophyllan a helical conformation by itself is not decisive and other factors are more important. Thus, in a single helical conformation schizophyllan stimulated TNF- $\alpha$  activity in U937, Th1, and human PBMC cells, but the native triple helical form was ineffective. A  $\beta$ -(1 $\rightarrow$ 3)(1 $\rightarrow$ 6)-glucan from *Glomerella cingulata* was effective against tumour growth in mice, regardless of whether it

existed as a triple helix, or in a single-helical conformation (Gomaa et al. 1992). Furthermore, chemically synthesized carbon 4- and 5- $\beta$ -linked oligosaccharides possess a high biological activity in the absence of any triple helical structure (Jamois et al. 2005).

Further study is required before we fully understand possible structure-function relationships in  $\beta$ -glucans. As already mentioned, orally administered  $\beta$ -glucans may be chemically and/or enzymatically modified following their ingestion, to generate smaller oligosaccharides (Hong et al. 2004). Thus, the actual molecules binding to the cell receptors to produce the effects discussed here may in fact be these smaller oligosaccharides. If this is the case, it may be possible, by careful choice of  $\beta$ -(1 $\rightarrow$ 3)(1 $\rightarrow$ 6)-endoglucanases or partial acid hydrolysis, to deliberately generate oligosaccharides of desired properties from chemically pure  $\beta$ -glucans. Consequently, assessing their biological activities could be fruitful. Unfortunately when enzymatic digestion of the structurally heterogeneous  $\beta$ -glucan from *Volvariella volvacea* generated several oligosaccharides with different branching frequencies, their individual anti-tumour activities were not reported (Kishida et al. 1989). In addition, it may be possible by selecting different cultural conditions to persuade the fungus to synthesize glucans with different branching frequencies, as reported for *Botryosphaera* spp. (Steluti et al. 2004), which might then be used as substrates for possible enzymatic modification. It is not known if the same approach would succeed with other  $\beta$ -glucan-producing fungi, although there may eventually be other culture-dependent methods to generate tailor-made  $\beta$ -glucans for work of this kind, once their biosynthesis and methods of assembly are better understood.

#### Adverse effects and efficient usage

When purified fungal  $\beta$ -glucans have been administered orally for treatment of human diseases, no adverse effects in humans have been recorded. However, co-administration of zymosan, sonifilan, grofolan, SSG, and non-steroidal anti-inflammatory drugs such as indomethacin has been reported to cause death (Yoshioka et al. 1998; Takahashi et al. 2001). Furthermore, particulate or slightly soluble fungal  $\beta$ -glucans given intravenously were reported to cause granuloma formation, microembolization, inflammation and pain (Di Luzio et al. 1979; Chesterman et al. 1981; Zekovic et al. 2005). Crude preparations from basidiocarps with high  $\beta$ -glucan levels may also have adverse side effects. Some wild mushrooms may contain high concentrations of toxic metals including arsenic, lead, cadmium and mercury (Borchers et al. 2004), and preparations with high protein contents may cause allergic responses including asthma (Borchers et al. 2004). Thus, it is important to restrict those used for human ingestion to approved and certified  $\beta$ -glucan products of known chemical composition.

In addition, although  $\beta$ -glucans are found in all fungi, they are complexed in cellular structures with other components, which can make them difficult to digest and be absorbed. Ideally those fungi producing exocellular  $\beta$ -glucans under carefully controlled laboratory conditions are more attractive sources and should be given specific attention, especially for

large-scale production. Furthermore the range of fungi reported to produce them continues to increase, and often the growth conditions known to stimulate their production are documented (Seviour et al. 1992; Selbmann et al. 2004; Steluti et al. 2004; Rezende et al. 2005; Leung et al. 2006).

## Conclusions

In summary, fungal  $\beta$ -glucans appear to be beneficial to humans with impaired immune systems, and those suffering from infectious diseases and cancer, as well as in helping patient recovery from chemotherapy and radiotherapy. They are also especially beneficial to middle-aged people, people with active and stressful lifestyles, and athletes. The mechanisms of action of these fungal  $\beta$ -glucans appear to depend on their capabilities to bind to cell receptors, which are known to include dectin-1, CR3, LacCer, and scavenger receptors. This event then leads to activation of multiple signal pathways which in turn promote immune responses in the affected cells. Data suggest that different fungal  $\beta$ -glucans have different effectiveness probably through their different binding affinity to each receptor, although the chemical purity of the glucans used in these studies is not always known. Correspondingly, the choice of which  $\beta$ -glucan to use may be important. Unfortunately, we still do not understand what structural features are best for inducing activities.

Future studies now need to characterize mechanistically how individual  $\beta$ -glucans of known structure bind to each receptor. Only then can we rationally improve this technology. This information may be obtained from 1) more precise identification of  $\beta$ -glucan structural diversity from the purification, characterization and screening of additional fungal  $\beta$ -glucans with structurally attractive overall features 2) synthesis of  $\beta$ -glucans with these optimal structural features by deliberate microbial engineering or by selection of appropriate culture conditions.

## REFERENCES

- Abel G, Czop JK, 1992. Stimulation of human monocyte  $\beta$ -glucan receptors by glucan particles induces production of TNF- $\alpha$  and IL-1  $\beta$ . *International Journal of Immunopharmacology* 14: 1363-1373.
- Acton SL, Scherer PE, Lodish HF, Krieger M, 1994. Expression cloning of SR-BI, a CD36-related class B scavenger receptor. *Journal of Biological Chemistry* 269: 21003-21009.
- Adachi Y, Ishii T, Ikeda Y, Hoshino A, Tamura H, Aketagawa J, Tanaka S, Ohno N, 2004. Characterization of  $\beta$ -glucan recognition site on C-type lectin, dectin1. *Infection and Immunity* 72: 4159-4171.
- Allendorf DJ, Yan J, Ross GD, Hansen RD, Baran JT, Subbarao K, Wang L, Haribabu B, 2005. C5a-mediated leukotriene B<sub>4</sub>-amplified neutrophil chemotaxis is essential in tumor immunotherapy facilitated by anti-tumor monoclonal antibody and  $\beta$ -glucan. *Journal of Immunology* 174: 7050-7056.
- Ara Y, Saito T, Takagi T, Hagiwara E, Miyagi Y, Sugiyama M, Kawamoto S, Ishii N, Yoshida T, Hanashi D, Koshino T, Okada H, Okuda K, 2001. Zymosan enhances the immune response to DNA vaccine for human immunodeficiency virus type-1 through the activation of complement system. *Immunology* 103: 98-105.
- Ariizumi K, Shen GL, Shikano S, Xu S, Ritter 3rd R, Kumamoto T, Edelbaum D, Morita A, Bergstresser PR, Takashima A, 2000. Identification of a novel, dendritic cell-associated molecule, dectin-1, by subtractive cDNA cloning. *Journal of Biological Chemistry* 275: 20157-20167.
- Arruda C, Franco MF, Kashino SS, Nascimento FR, Fazioli Rdos A, Vaz CA, Russo M, Calich VL, 2002. Interleukin-12 protects mice against disseminated infection caused by *Paracoccidioides brasiliensis* but enhances pulmonary inflammation. *Clinical Immunology* 103: 185-195.
- Assanasen C, Mineo C, Seetharam D, Yuhanna IS, Marcel YL, Connelly MA, Williams DL, de la Llera-Moya M, Shaul PW, Silver DL, 2005. Cholesterol binding, efflux, and a PDZ-interacting domain of scavenger receptor-BI mediate HDL-initiated signaling. *Journal of Clinical Investigation* 115: 969-977.
- Berner MD, Sura ME, Alves BN, Hunter KW Jr, 2005. IFN- $\gamma$  primes macrophages for enhanced TNF- $\alpha$  expression in response to stimulatory and non-stimulatory amounts of microparticulate  $\beta$ -glucan. *Immunology Letters* 98: 115-122.
- Beutler B, Cerami A, 1989. The biology of cachectin/TNF—a primary mediator of the host response. *Annual Review of Immunology* 7: 625-655.
- Borchers AT, Keen CL, Gershwin ME, 2004. Mushrooms, tumors, and immunity: an update. *Experimental Biology and Medicine (Maywood)* 229: 393-406.
- Brown GD, 2006. Dectin-1: a signalling non-TLR pattern-recognition receptor. *Nature Reviews Immunology* 6: 33-43.
- Brown GD, Gordon S, 2001. Immune recognition. A new receptor for  $\beta$ -glucans. *Nature* 413: 36-37.
- Brown GD, Gordon S, 2005. Immune recognition of fungal  $\beta$ -glucans. *Cellular Microbiology* 7: 471-479.
- Brown GD, Taylor PR, Reid DM, Willment JA, Williams DL, Martínez-Pomares L, Wong SY, Gordon S, 2002. Dectin-1 is a major  $\beta$ -glucan receptor on macrophages. *Journal of Experimental Medicine* 196: 407-412.
- Calich VL, Vaz CA, Burger E, 1998. Immunity to *Paracoccidioides brasiliensis* infection. *Research in Immunology* 149: 407-417 discussion 499-500.
- Chesterman H, Heywood R, Allen TR, Street AE, Edmondson NA, Prentice DE, 1981. The intravenous toxicity of lentinan to the beagle dog. *Toxicology Letters* 9: 87-90.
- Cheung NK, Modak S, 2002. Oral (1 $\rightarrow$ 3),(1 $\rightarrow$ 4)- $\beta$ -D-glucan synergizes with antiganglioside GD2 monoclonal antibody 3F8 in the therapy of neuroblastoma. *Clinical Cancer Research* 8: 1217-1223.
- Cheung NK, Modak S, Vickers A, Knuckles B, 2002. Orally administered  $\beta$ -glucans enhance anti-tumor effects of monoclonal antibodies. *Cancer Immunology, Immunotherapy* 51: 557-564.
- de la Rosa G, Yanez-Mo M, Samaneigo R, Serrano-Gomez D, Martínez-Munoz L, Fernandez-Ruiz E, Longo N, Sanchez-Madrid F, Corbi AL, Sanchez-Mateos P, 2005. Regulated recruitment of DC-SIGN to cell-cell contact regions during zymosan-induced human dendritic cell aggregation. *Journal of Leukocyte Biology* 77: 699-709.
- Demleitner S, Kraus J, Franz G, 1992. Synthesis and anti-tumour activity of sulfoalkyl derivatives of curdian and lichenan. *Carbohydrate Research* 226: 247-252.
- Di Luzio NR, Williams DL, McNamee RB, Edwards BF, Kitahama A, 1979. Comparative tumor-inhibitory and anti-bacterial activity of soluble and particulate glucan. *International Journal of Cancer* 24: 773-779.
- Di Luzio NR, Williams DL, McNamee RB, Malshet VG, 1980. Comparative evaluation of the tumor inhibitory and antibacterial activity of solubilized and particulate glucan. *Recent Results in Cancer Research* 75: 165-172.
- Diniz SN, Nomizo R, Cisalpino PS, Teixeira MM, Brown GD, Mantovani A, Gordon S, Reis LF, Dias AA, 2004. PTX3 function

- as an opsonin for the dectin-1-dependent internalization of zymosan by macrophages. *Journal of Leukocyte Biology* 75: 649–656.
- Drandarska J, Kussovski V, Nikolaeva S, Markova N, 2005. Combined immunomodulating effects of BCG and Lentinan after intranasal application in guinea pigs. *International Immunopharmacology* 5: 795–803.
- Du Z, Kelly E, Mecklenbrauker J, Agle L, Herrero C, Paik P, Ivashkiv LB, 2006. Selective regulation of IL-10 signaling and function by zymosan. *Journal of Immunology* 176: 4785–4792.
- Duenas AI, Aceves M, Orduna A, Diaz R, Sanchez Crespo M, Garcia-Rodriguez C, 2006. *Francisella tularensis* LPS induces the production of cytokines in human monocytes and signals via toll-like receptor 4 with much lower potency than *E. coli* LPS. *International Immunology* 18: 785–795.
- Engering A, Geijtenbeek TB, van Vliet SJ, Wijers M, van Liempt E, Demareux N, Lanzavecchia A, Fransen J, Figdor CG, Piguet V, van Kooyk Y, 2002. The dendritic cell-specific adhesion receptor DC-SIGN internalizes antigen for presentation to T cells. *Journal of Immunology* 168: 2118–2126.
- Evans SE, Hahn PY, McCann F, Kottom TJ, Pavlovic ZV, Limper AH, 2005. *Pneumocystis* cell wall  $\beta$ -glucans stimulate alveolar epithelial cell chemokine generation through nuclear factor- $\kappa$ B-dependent mechanisms. *American Journal of Respiratory Cellular and Molecular Biology* 32: 490–497.
- Falch BH, Espevik T, Ryan L, Stokke BT, 2000. The cytokine stimulating activity of (1 $\rightarrow$ 3)- $\beta$ -D-glucans is dependent on the triple helix conformation. *Carbohydrate Research* 329: 587–596.
- Gantner BN, Simmons RM, Canavera SJ, Akira S, Underhill DM, 2003. Collaborative induction of inflammatory responses by dectin-1 and toll-like receptor 2. *Journal of Experimental Medicine* 197: 1107–1117.
- Gantner BN, Simmons RM, Underhill DM, 2005. Dectin-1 mediates macrophage recognition of *Candida albicans* yeast but not filaments. *The EMBO Journal* 24: 1277–1286.
- Garcia-Lora A, Martinez M, Pedrinaci S, Garrido F, 2003. Different regulation of PKC isoenzymes and MAPK by PSK and IL-2 in the proliferative and cytotoxic activities of the NKL human natural killer cell line. *Cancer Immunology, Immunotherapy* 52: 59–64.
- Garcia-Lora A, Pedrinaci S, Garrido F, 2001. Protein-bound polysaccharide K and interleukin-2 regulate different nuclear transcription factors in the NKL human natural killer cell line. *Cancer Immunology, Immunotherapy*: CII 50: 191–198.
- Gomaa K, Kraus J, Roskopf F, Roper H, Franz G, 1992. Antitumour and immunological activity of a  $\beta$  1 $\rightarrow$ 3/1 $\rightarrow$ 6 glucan from *Glomerella cingulata*. *Journal of Cancer Research and Clinical Oncology* 118: 136–140.
- Gordon M, Bihari B, Goosby E, Gorter R, Greco M, Guralnik M, Mimura T, Rudnicki V, Wong R, Kaneko Y, 1998. A placebo-controlled trial of the immune modulator, lentinan, in HIV-positive patients: a phase I/II trial. *Journal of Medicine* 29: 305–330.
- Gordon M, Guralnik M, Kaneko Y, Mimura T, Goodgame J, DeMarzo C, Pierce D, Baker M, Lang W, 1995. A phase II controlled study of a combination of the immune modulator, lentinan, with didanosine (ddi) in HIV patients with CD4 cells of 200–500 mm<sup>-3</sup>. *Journal of Medicine* 26: 193–207.
- Greenwald P, Clifford CK, Milner JA, 2001. Diet and cancer prevention. *European Journal of Cancer* 37: 948–965.
- Gross O, Gewies A, Finger K, Schafer M, Sparwasser T, Peschel C, Forster I, Ruland J, 2006. Card9 controls a non-TLR signalling pathway for innate anti-fungal immunity. *Nature* 442: 651–656.
- Grunebach F, Weck MM, Reichert J, Brossart P, 2002. Molecular and functional characterization of human dectin-1. *Experimental Hematology* 30: 1309–1315.
- Gu YH, Takagi Y, Nakamura T, Hasegawa T, Suzuki I, Oshima M, Tawaraya H, Niwano Y, 2005. Enhancement of radioprotection and anti-tumor immunity by yeast-derived  $\beta$ -glucan in mice. *Journal of Medical Food* 8: 154–158.
- Hagiwara SY, Takahashi M, Shen Y, Kaihou S, Tomiyama T, Yazawa M, Tamai Y, Sin Y, Kazusaka A, Terazawa M, 2005. A phytochemical in the edible Tamogi-take mushroom (*Pleurotus cornucopiae*), D-mannitol, inhibits ACE activity and lowers the blood pressure of spontaneously hypertensive rats. *Bioscience, Biotechnology and Biochemistry* 69: 1603–1605.
- Hahn PY, Evans SE, Kottom TJ, Standing JE, Pagano RE, Limper AH, 2003. *Pneumocystis carinii* cell wall  $\beta$ -glucan induces release of macrophage inflammatory protein-2 from alveolar epithelial cells via a lactosylceramide-mediated mechanism. *Journal of Biological Chemistry* 278: 2043–2050.
- Harada T, Miura NN, Adachi Y, Nakajima M, Yadomae T, Ohno N, 2002. IFN- $\gamma$  induction by SCG, 1,3- $\beta$ -D-glucan from *Sparassis crispa*, in DBA/2 mice in vitro. *Journal of Interferon and Cytokine Research* 22: 1227–1239.
- Harrison LM, van Haaften WC, Tesh VL, 2004. Regulation of proinflammatory cytokine expression by Shiga toxin 1 and/or lipopolysaccharides in the human monocytic cell line THP-1. *Infection and Immunity* 72: 2618–2627.
- Hernanz-Falcon P, Arce I, Roda-Navarro P, Fernandez-Ruiz E, 2001. Cloning of human DECTIN-1, a novel C-type lectin-like receptor gene expressed on dendritic cells. *Immunogenetics* 53: 288–295.
- Herre J, Gordon S, Brown GD, 2004a. Dectin-1 and its role in the recognition of  $\beta$ -glucans by macrophages. *Molecular Immunology* 40: 869–876.
- Herre J, Willment JA, Gordon S, Brown GD, 2004b. The role of dectin-1 in antifungal immunity. *Critical Review of Immunology* 24: 193–203.
- Hetland G, Ohno N, Aaberge IS, Lovik M, 2000. Protective effect of  $\beta$ -glucan against systemic *Streptococcus pneumoniae* infection in mice. *FEMS Immunology and Medical Microbiology* 27: 111–116.
- Hobson RP, Munro CA, Bates S, MacCallum DM, Cutler JE, Heinsbroek SE, Brown GD, Odds FC, Gow NA, 2004. Loss of cell wall mannosylphosphate in *Candida albicans* does not influence macrophage recognition. *Journal of Biological Chemistry* 279: 39628–39635.
- Hohl TM, Van Epps HL, Rivera A, Morgan LA, Chen PL, Feldmesser M, Pamer EG, 2005. *Aspergillus fumigatus* triggers inflammatory responses by stage-specific  $\beta$ -glucan display. *PLoS Pathogens* 1: e30.
- Hong F, Yan J, Baran JT, Allendorf DJ, Hansen RD, Ostroff GR, Xing PX, Cheung NK, Ross GD, 2004. Mechanism by which orally administered  $\beta$ -1,3-glucans enhance the tumoricidal activity of antitumor monoclonal antibodies in murine tumor models. *Journal of Immunology* 173: 797–806.
- Ito K, Nakazato H, Koike A, Takagi H, Saji S, Baba S, Mai M, Sakamoto J, Ohashi Y, 2004. Long-term effect of 5-fluorouracil enhanced by intermittent administration of polysaccharide K after curative resection of colon cancer. A randomized controlled trial for 7-year follow-up. *International Journal of Colorectal Diseases* 19: 157–164.
- Iwabuchi K, Nagaoka I, 2002. Lactosylceramide-enriched glycosphingolipid signaling domain mediates superoxide generation from human neutrophils. *Blood* 100: 1454–1464.
- Jamois F, Ferrieres V, Guegan JP, Yvin JC, Plusquellec D, Vetricka V, 2005. Glucan-like synthetic oligosaccharides: iterative synthesis of linear oligo- $\beta$ -(1,3)-glucans and immunostimulatory effects. *Glycobiology* 15: 393–407.
- Jayus, McDougall BM, Seviour RJ, 2004. Purification and characterization of the (1 $\rightarrow$ 3)- $\beta$ -glucanases from *Acremonium* sp. IMI 383068. *FEMS Microbiological Letters* 230: 259–264.

- Johnson JJ, Kirkwood A, Misaki A, Nelson T, Scalettie J, Smith F, 1963. Structure of a new glucan. *Chemistry Industry. (London)* 41: 820-822.
- Jung K, Ha Y, Ha SK, Han DU, Kim DW, Moon WK Chae C, 2004. Antiviral effect of *Saccharomyces cerevisiae*  $\beta$ -glucan to swine influenza virus by increased production of interferon-gamma and nitric oxide. *Journal of Veterinary Medicine Series B Infectious Diseases and Veterinary Public Health* 51: 72-76.
- Kabir Y, Kimura S, 1989. Dietary mushrooms reduce blood pressure in spontaneously hypertensive rats (SHR). *Journal of Nutritional Science and Vitaminology (Tokyo)* 35: 91-94.
- Kabir Y, Yamaguchi M, Kimura S, 1987. Effect of shiitake (*Lentinus edodes*) and maitake (*Grifola frondosa*) mushrooms on blood pressure and plasma lipids of spontaneously hypertensive rats. *Journal of Nutritional Science and Vitaminology (Tokyo)* 33: 341-346.
- Kaiser AB, Kernodle DS, 1998. Synergism between poly-(1-6)- $\beta$ -D-glucopyranosyl-(1-3)- $\beta$ -D-glucopyranose glucan and cefazolin in prophylaxis of staphylococcal wound infection in a guinea pig model. *Antimicrobial Agents and Chemotherapy* 42: 2449-2451.
- Kakumu S, Ishikawa T, Wakita T, Yoshioka K, Ito Y, Shinagawa T, 1991. Effect of sizofiran, a polysaccharide, on interferon gamma, antibody production and lymphocyte proliferation specific for hepatitis B virus antigen in patients with chronic hepatitis B. *International Journal of Immunopharmacology* 13: 969-975.
- Kanazawa M, Mori Y, Yoshihara K, Iwadata M, Suzuki S, Endoh Y, Ohki S, Takita K, Sekikawa K, Takenoshita S, 2004. Effect of PSK on the maturation of dendritic cells derived from human peripheral blood monocytes. *Immunological Letters* 91: 229-238.
- Kanse SM, Matz RL, Preissner KT, Peter K, 2004. Promotion of leukocyte adhesion by a novel interaction between vitronectin and the  $\beta$ 2 integrin Mac-1 ( $\alpha$ M $\beta$ 2, CD11b/CD18). *Arteriosclerosis, Thrombosis, and Vascular Biology* 24: 2251-2256.
- Kataoka K, Muta T, Yamazaki S, Takeshige K, 2002. Activation of macrophages by linear (1 $\rightarrow$ 3)- $\beta$ -D-glucans. Implications for the recognition of fungi by innate immunity. *Journal of Biological Chemistry* 277: 36825-36831.
- Kawai T, Takeuchi O, Fujita T, Inoue J, Muhlradt PF, Sato S, Hoshino K, Akira S, 2001. Lipopolysaccharide stimulates the MyD88-independent pathway and results in activation of IFN-regulatory factor 3 and the expression of a subset of lipopolysaccharide-inducible genes. *Journal of Immunology* 167: 5887-5894.
- Kernodle DS, Gates H, Kaiser AB, 1998. Prophylactic anti-infective activity of poly-[1-6]- $\beta$ -D-glucopyranosyl-[1-3]- $\beta$ -D-glucopyranose glucan in a guinea pig model of staphylococcal wound infection. *Antimicrobial Agents and Chemotherapy* 42: 545-549.
- Kiho T, Kobayashi T, Morimoto H, Usui S, Ukai S, Hirano K, Aizawa K, Inakuma T, 2000. Structural features of an anti-diabetic polysaccharide (TAP) from *Tremella aurantia*. *Chemical and Pharmaceutical Bulletin (Tokyo)* 48: 1793-1795.
- Kiho T, Matsushita M, Usui S, Ukai S, 1998. Biological activities of (1 $\rightarrow$ 3)- $\beta$ -D-glucans with reducing glucose side chains. *Bioscience, Biotechnology and Biochemistry* 62: 570-572.
- Kiho T, Morimoto H, Sakushima M, Usui S, Ukai S, 1995. Polysaccharides in fungi. XXXV. Anti diabetic activity of an acidic polysaccharide from the fruiting bodies of *Tremella aurantia*. *Biological and Pharmaceutical Bulletin* 18: 1627-1629.
- Kikumoto SMT, Kimura K, 1971. Polysaccharide produced by *Schizophyllum commune*. Part II. Chemical structure of an extracellular polysaccharide. *Journal of Agricultural and Chemical Society Japan* 45: 162-168.
- Kim YW, Kim KH, Choi HJ, Lee DS, 2005. Anti-diabetic activity of  $\beta$ -glucans and their enzymatically hydrolyzed oligosaccharides from *Agaricus blazei*. *Biotechnological Letters* 27: 483-487.
- Kishida E, Sone Y, Misaki A, 1989. Purification of an antitumor-active, branched (1 $\rightarrow$ 3)- $\beta$ -D-glucan from *Volvariella volvacea*, and elucidation of its fine structure. *Carbohydrate Research* 193: 227-239.
- Klis FM, de Groot P, Hellingwerf K, 2001. Molecular organization of the cell wall of *Candida albicans*. *Medical Mycology* 39: 1-8.
- Kobayashi H, Matsunaga K, Oguchi Y, 1995. Antimetastatic effects of PSK (Krestin), a protein-bound polysaccharide obtained from *basidiomycetes*: an overview. *Cancer Epidemiology, Biomarkers and Prevention* 4: 275-281.
- Kobayashi H, Yoshida R, Kanada Y, Fukuda Y, Yagyu T, Inagaki K, Kondo T, Kurita N, Suzuki M, Kanayama N, Terao T, 2005. Suppressing effects of daily oral supplementation of  $\beta$ -glucan extracted from *Agaricus blazei* Murill on spontaneous and peritoneal disseminated metastasis in mouse model. *Journal of Cancer Research and Clinical Oncology* 131: 527-538.
- Kournikakis B, Mandeville R, Brousseau P, Ostroff G, 2003. Anthrax-protective effects of yeast  $\beta$  1,3 glucans. *Medscape General Medicine* 5: 1.
- Kukan M, Szatmary Z, Lutterova M, Kuba D, Vajdova K, Horecky J, 2004. Effects of sizofiran on endotoxin-enhanced cold ischemia-reperfusion injury of the rat liver. *Physiological Research* 53: 431-437.
- Kurashige S, Akuzawa Y, Endo F, 1997. Effects of *Lentinus edodes*, *Grifola frondosa* and *Pleurotus ostreatus* administration on cancer outbreak, and activities of macrophages and lymphocytes in mice treated with a carcinogen, N-butyl-N-butanoinitrosamine. *Immunopharmacology and Immunotoxicology* 19: 175-183.
- Ladanyi A, Timar J, Lapis K, 1993. Effect of lentinan on macrophage cytotoxicity against metastatic tumor cells. *Cancer Immunology and Immunotherapy* 36: 123-126.
- Lapaque N, Takeuchi O, Corrales F, Akira S, Moriyon I, Howard JC, Gorvel JP, 2006. Differential inductions of TNF- $\alpha$  and iGTP, iIGP by structurally diverse classic and non-classic lipopolysaccharides. *Cell Microbiology* 8: 401-413.
- Lebron F, Vassallo R, Puri V, Limper AH, 2003. *Pneumocystis carinii* cell wall  $\beta$ -glucans initiate macrophage inflammatory responses through NF-kappaB activation. *Journal of Biological Chemistry* 278: 25001-25008.
- Leung PH, Zhang QX, Wu JY, 2006. Mycelium cultivation, chemical composition and antitumor activity of a *Tolyposcladium* sp. fungus isolated from wild *Cordyceps sinensis*. *Journal of Applied Microbiology* 101: 275-283.
- Li B, Allendorf DJ, Hansen R, Marroquin J, Ding C, Cramer DE, Yan J, 2006. Yeast  $\beta$ -glucan amplifies phagocyte killing of iC3b-opsonized tumor cells via complement receptor 3-Syk-phosphatidylinositol 3-kinase pathway. *Journal of Immunology* 177: 1661-1669.
- Liang J, Melican D, Cafro L, Palace G, Fisetto L, Armstrong R, Patchen ML, 1998. Enhanced clearance of a multiple antibiotic resistant *Staphylococcus aureus* in rats treated with PGG-glucan is associated with increased leukocyte counts and increased neutrophil oxidative burst activity. *International Journal of Immunopharmacology* 20: 595-614.
- Lindequist U, Niedermeyer TH, Julich WD, 2005. The pharmacological potential of mushrooms. *Evidence Based Complementary and Alternative Medicine* 2: 285-299.
- Lo HC, Tsai FA, Wasser SP, Yang JG, Huang BM, 2006. Effects of ingested fruiting bodies, submerged culture biomass, and acidic polysaccharide glucuronoxylomannan of *Tremella mesenterica* Retz.:Fr. on glycemic responses in normal and diabetic rats. *Life Science* 78: 1957-1966.
- Lucas EH, Byerrum RU, Clarke DA, Reilly HC, Stevens JA, Stock CC, 1958. Production of oncostatic principles *in vivo* and *in vitro* by species of the genus *Calvatia*. *Antibiotics Annual* 6: 493-496.
- Mantegazza AR, Barrio MM, Moutel S, Bover I, Weck M, Brossart P, Teillaud JL, Mordoh J, 2004. CD63 tetraspanin slows down cell

- migration and translocates to the endosomal-lysosomal-MHCs route after extracellular stimuli in human immature dendritic cells. *Blood* 104: 1183-1190.
- Markova N, Kussovski V, Drandarska I, Nikolaeva S, Georgieva N, Radoucheva T, 2003. Protective activity of Lentinan in experimental tuberculosis. *International Immunopharmacology* 3: 1557-1562.
- Markova N, Michailova L, Kussovski V, Jourdanova M, Radoucheva T, 2005. Intranasal application of lentinan enhances bactericidal activity of rat alveolar macrophages against *Mycobacterium tuberculosis*. *Pharmazie* 60: 42-48.
- Marshall AS, Willment JA, Lin HH, Williams DL, Gordon S, Brown GD, 2004. Identification and characterization of a novel human myeloid inhibitory C-type lectin-like receptor (MIEL) that is predominantly expressed on granulocytes and monocytes. *Journal of Biological Chemistry* 279: 14792-14802.
- Martin KL, McDougall BM, Unkles SE, Seviour RJ, 2006. The three  $\beta$ -1,3-glucanases from *Acremonium blochii* strain C59 appear to be encoded by separate genes. *Mycological Research* 110: 66-74.
- Mayell M, 2001. Maitake extracts and their therapeutic potential. *Alternative Medicine Review* 6: 48-60.
- McIntosh M, Stone BA, Stanisich VA, 2005. Curdlan and other bacterial (1 $\rightarrow$ 3)- $\beta$ -D-glucans. *Applied Microbiology and Biotechnology* 68: 163-173.
- Mineo C, Yuhanna IS, Quon MJ, Shaul FW, 2003. High density lipoprotein-induced endothelial nitric-oxide synthase activation is mediated by Akt and MAP kinases. *Journal of Biological Chemistry* 278: 9142-9149.
- Misaki A, Kakuta M, Sasaki T, Tanaka M, Miyaji H, 1981. Studies on interrelation of structure and antitumor effects of polysaccharides: antitumor action of periodate-modified, branched (1 $\rightarrow$ 3)- $\beta$ -D-glucan of *Auricularia auricula-judae*, and other polysaccharides containing (1 $\rightarrow$ 3)-glycosidic linkages. *Carbohydrate Research* 92: 115-129.
- Misaki A, Kawaguchi K, Miyaji H, Nagae H, Hokkoku S, Kakuta M, Sasaki T, 1984. Structure of pestalotan, a highly branched (1 $\rightarrow$ 3)- $\beta$ -D-glucan elaborated by *Pestalotia* sp. 815, and the enhancement of its antitumor activity by polyol modification of the side chains. *Carbohydrate Research* 129: 209-227.
- Mitamura T, Sakamoto S, Suzuki S, Yoshimura S, Maemura M, Kudo H, 2000. Effects of lentinan on colorectal carcinogenesis in mice with ulcerative colitis. *Oncology Reports* 7: 599-601.
- Miyazaki T, Oikawa N, Yadomae T, Yamada H, Yamada Y, Hsu HY, Ito H, 1979. Relationship between the chemical structure and anti-tumour activity of glucans prepared from *Grifora umbellata*. *Carbohydrate Research* 69: 165-170.
- Mizuno M, Minato K, Ito H, Kawade M, Terai H, Tsuchida H, 1999. Anti-tumor polysaccharide from the mycelium of liquid-cultured *Agaricus blazei* mill. *Biochemistry and Molecular Biology International* 47: 707-714.
- Modak S, Koehne G, Vickers A, O'Reilly RJ, Cheung NK, 2005. Rituximab therapy of lymphoma is enhanced by orally administered (1 $\rightarrow$ 3),(1 $\rightarrow$ 4)-D- $\beta$ -glucan. *Leukaemia Research* 29: 679-683.
- Mueller A, Raptis J, Rice PJ, Kalbfleisch JH, Stout RD, Ensley HE, Browder W, Williams DL, 2000. The influence of glucan polymer structure and solution conformation on binding to (1 $\rightarrow$ 3)- $\beta$ -D-glucan receptors in a human monocyte-like cell line. *Glycobiology* 10: 339-346.
- Munemoto Y, Iida Y, Abe J, Saito H, Fujisawa K, Kasahara Y, Mitsui T, Asada Y, Miura S, 2002. Significance of postoperative adjuvant immunotherapy after curative resection of colorectal cancers: association between host or tumor factors and survival. *International Journal of Oncology* 20: 403-411.
- Munz C, Steinman RM, Fujii S, 2005. Dendritic cell maturation by innate lymphocytes: coordinated stimulation of innate and adaptive immunity. *Journal of Experimental Medicine* 202: 203-207.
- Mushiake H, Tsunoda T, Nukatsuka M, Shima K, Fukushima M, Tahara H, 2005. Dendritic cells might be one of key factors for eliciting antitumor effect by chemoimmunotherapy *in vivo*. *Cancer Immunology and Immunotherapy* 54: 120-128.
- Nakano H, Namatame K, Nemoto H, Motohashi H, Nishiyama K, Kumada K, 1999. A multi-institutional prospective study of lentinan in advanced gastric cancer patients with unresectable and recurrent diseases: effect on prolongation of survival and improvement of quality of life. Kanagawa Lentinan Research Group. *Hepatogastroenterology* 46: 2662-2668.
- Nakazato H, Koike A, Saji S, Ogawa N, Sakamoto J, 1994. Efficacy of immunochemotherapy as adjuvant treatment after curative resection of gastric cancer. Study Group of Immunochemotherapy with PSK for Gastric Cancer. *Lancet* 343: 1122-1126.
- Nam SW, Park JY, Ramasamy A, Shevade S, Islam A, Long PM, Park CK, Park SE, Kim SY, Lee SH, Park WS, Yoo NJ, Liu ET, Miller LD, Lee JY, 2005. Molecular changes from dysplastic nodule to hepatocellular carcinoma through gene expression profiling. *Hepatology* 42: 809-818.
- Netea MG, Gow NA, Munro CA, Bates S, Collins C, Ferwerda G, Hobson RP, Bertram G, Hughes HB, Jansen T, Jacobs L, Buurman ET, Gijzen K, Williams DL, Torensma R, McKinnon A, MacCallum DM, Odds FC, Van der Meer JW, Brown AJ, Kullberg BJ, 2006. Immune sensing of *Candida albicans* requires cooperative recognition of mannans and glucans by lectin and toll-like receptors. *Journal of Clinical Investigation* 116: 1642-1650.
- Nicolosi R, Belli SJ, Bistran BR, Greenberg I, Forse RA, Blackburn GL, 1999. Plasma lipid changes after supplementation with  $\beta$ -glucan fiber from yeast. *American Journal of Clinical Nutrition* 70: 208-212.
- Nilsson S, Norlen BJ, Widmark A, 2004. A systematic overview of radiation therapy effects in prostate cancer. *Acta Oncology* 43: 316-381.
- Ohwada S, Ikeya T, Yokomori T, Kusaba T, Roppongi T, Takahashi T, Nakamura S, Kakinuma S, Iwazaki S, Ishikawa H, Kawate S, Nakajima T, Morishita Y, 2004. Adjuvant immunotherapy with oral tegafur/uracil plus PSK in patients with stage II or III colorectal cancer: a randomised controlled study. *British Journal of Cancer* 90: 1003-1010.
- Okamoto T, Kodoi R, Nonaka Y, Fukuda I, Hashimoto T, Kanazawa K, Mizuno M, Ashida H, 2004. Lentinan from shiitake mushroom (*Lentinus edodes*) suppresses expression of cytochrome P450 1A subfamily in the mouse liver. *Biofactors* 21: 407-409.
- Ozinsky A, Underhill DM, Fontenot JD, Hajjar AM, Smith KD, Wilson CB, Schroeder L, Aderem A, 2000. The repertoire for pattern recognition of pathogens by the innate immune system is defined by cooperation between toll-like receptors. *Proceedings of the National Academy of Sciences USA* 97: 13766-13771.
- Pajma AS, Feizi T, Zhang Y, Stoll MS, Lawson AM, Diaz-Rodriguez E, Campanero-Rhodes MA, Costa J, Gordon S, Brown GD, Chai W, 2006. Ligands for the  $\beta$ -glucan receptor, Dectin-1, assigned using 'designer' microarrays of oligosaccharide probes (neoglycolipids) generated from glucan polysaccharides. *Journal of Biological Chemistry* 281: 5771-5779.
- Pelizon AC, Kaneno R, Soares AM, Meira DA, Sartori A, 2005. Immunomodulatory activities associated with  $\beta$ -glucan derived from *Saccharomyces cerevisiae*. *Physiological Research* 54: 557-564.
- Pitson SM, Seviour RJ, McDougall BM, 1996a. Proteolytic inactivation of an extracellular (1 $\rightarrow$ 3)- $\beta$ -glucanase from the fungus *Acremonium persicinum* is associated with growth at neutral or alkaline medium pH. *FEMS Microbiology Letters* 145: 287-293.

- Pitson SM, Seviour RJ, McDougall BM, Stone BA, Sadek M, 1996b. Purification and characterization of an extracellular (1→6)-β-glucanase from the filamentous fungus *Acremonium persicinum*. *Biochemical Journal* 316: 841–846.
- Portera CA, Love EJ, Memore L, Zhang L, Muller A, Browder W, Williams DL, 1997. Effect of macrophage stimulation on collagen biosynthesis in the healing wound. *American Surgery* 63: 125–131.
- Rezende MI, Barbosa AM, Vasconcelos AF, Haddad R, Dekker RF, 2005. Growth and production of laccases by the ligninolytic fungi, *Pleurotus ostreatus* and *Botryosphaeria rhodina*, cultured on basal medium containing the herbicide, Scepter (imazaquin). *Journal of Basic Microbiology* 45: 460–469.
- Rice PJ, Kelley JL, Kogan G, Ensley HE, Kalbfleisch JH, Browder IW, Williams DL, 2002. Human monocyte scavenger receptors are pattern recognition receptors for (1→3)-β-D-glucans. *Journal of Leukocyte Biology* 72: 140–146.
- Roeder A, Kirschning CJ, Rupec RA, Schaller M, Weindl G, Korting HC, 2004. Toll-like receptors as key mediators in innate antifungal immunity. *Medical Mycology* 42: 485–498.
- Rogers NC, Slack EC, Edwards AD, Nolte MA, Schulz O, Schweighoffer E, Williams DL, Gordon S, Tybulewicz VL, Brown GD, Reis e Sousa C, 2005. Syk-dependent cytokine induction by Dectin-1 reveals a novel pattern recognition pathway for C type lectins. *Immunity* 22: 507–517.
- Ross GD, 2000. Regulation of the adhesion versus cytotoxic functions of the Mac-1/CR3/αMβ2-integrin glycoprotein. *Critical Review of Immunology* 20: 197–222.
- Saijo S, Fujikado N, Furuta T, Chung SH, Kotaki H, Seki K, Sudo K, Akira S, Adachi Y, Ohno N, Kinjo T, Nakamura K, Kawakami K, Iwakura Y, 2007. Dectin-1 is required for host defense against *Pneumocystis carinii* but not against *Candida albicans*. *Nature Immunology* 8: 39–46.
- Sakamoto J, Morita S, Oba K, Matsui T, Kobayashi M, Nakazato H, Ohashi Y, 2006. Efficacy of adjuvant immunochemotherapy with polysaccharide K for patients with curatively resected colorectal cancer: a meta-analysis of centrally randomized controlled clinical trials. *Cancer Immunology and Immunotherapy* 55: 404–411.
- Sasaki T, Takasuka N, 1976. Further study of the structure of lentinan, an anti-tumor polysaccharide from *Lentinus edodes*. *Carbohydrate Research* 47: 99–104.
- Sato M, Sano H, Iwaki D, Kudo K, Konishi M, Takahashi H, Takahashi T, Imaizumi H, Asai Y, Kuroki Y, 2003. Direct binding of Toll-like receptor 2 to zymosan, and zymosan-induced NF-κB activation and TNF-α secretion are down-regulated by lung collectin surfactant protein A. *Journal of Immunology* 171: 417–425.
- Schmid F, Stone BA, McDougall BM, Bacic A, Martin KL, Brownlee RT, Chai E, Seviour RJ, 2001. Structure of epiglucan, a highly side-chain/branched (1→3;1→6)-β-glucan from the micro fungus *Epicoccum nigrum* Ehrenb. ex Schlecht. *Carbohydrate Research* 331: 163–171.
- Selbmann L, Crognale S, Petruccioli M, 2004. β-glucan production by *Botryosphaeria rhodina* in different bench-top bioreactors. *Journal of Applied Microbiology* 96: 1074–1081.
- Serrano-Gomez D, Dominguez-Soto A, Ancochea J, Jimenez-Heffernan JA, Leal JA, Corbi AL, 2004. Dendritic cell-specific intercellular adhesion molecule 3-grabbing nonintegrin mediates binding and internalization of *Aspergillus fumigatus* conidia by dendritic cells and macrophages. *Journal of Immunology* 173: 5635–5643.
- Seviour RJ, Stasinopoulos SJ, Auer DPF, 1992. Production of pululan and other exopolysaccharides by filamentous fungi. *Critical Reviews in Biotechnology* 12: 279–298.
- Sier CF, Gelderman KA, Prins FA, Gorter A, 2004. β-glucan enhanced killing of renal cell carcinoma micrometastases by monoclonal antibody G250 directed complement activation. *International Journal of Cancer* 109: 900–908.
- Sionov E, Mendlovic S, Segal E, 2005. Experimental systemic murine aspergillosis: treatment with polyene and caspofungin combination and G-CSF. *Journal of Antimicrobials and Chemotherapy* 56: 594–597.
- Sporn MB, Liby KT, 2005. Cancer chemoprevention: scientific promise, clinical uncertainty. *Nature Clinical Practice and Oncology* 2: 518–525.
- Steele C, Marrero L, Swain S, Harmsen AG, Zheng M, Brown GD, Gordon S, Shellito JE, Kolls JK, 2003. Alveolar macrophage-mediated killing of *Pneumocystis carinii* f. sp. muris involves molecular recognition by the dectin-1 β-glucan receptor. *Journal of Experimental Medicine* 198: 1677–1688.
- Steluti RM, Giese EC, Figgato MM, Sumiya AF, Covizzi LG, Job AE, Cardoso MS, Corradi da Silva Mde L, Dekker RF, Barbosa AM, 2004. Comparison of botryosphaeran production by the ascomyceteous fungus *Botryosphaeria* sp., grown on different carbohydrate carbon sources, and their partial structural features. *Journal of Basic Microbiology* 44: 480–486.
- Strong K, Mathers C, Leeder S, Beaglehole R, 2005. Preventing chronic diseases: how many lives can we save? *Lancet* 366: 1578–1582.
- Suram S, Brown GD, Ghosh M, Gordon S, Loper R, Taylor PR, Akira S, Uematsu S, Williams DL, Leslie CC, 2006. Regulation of cytosolic phospholipase A2 activation and cyclooxygenase 2 expression in macrophages by the β-glucan receptor. *Journal of Biological Chemistry* 281: 5506–5514.
- Suzuki Y, Adachi Y, Ohno N, Yadomae T, 2001. Th1/Th2-Balancing immunomodulating activity of gel-forming (1→3)-β-glucans from fungi. *Biological and Pharmaceutical Bulletin* 24: 811–819.
- Takahashi H, Ohno N, Adachi Y, Yadomae T, 2001. Association of immunological disorders in lethal side effect of NSAIDs on β-glucan-administered mice. *FEMS Immunology and Medical Microbiology* 31: 1–14.
- Takaku T, Kimura Y, Okuda H, 2001. Isolation of an antitumor compound from *Agaricus blazei* Murill and its mechanism of action. *Journal of Nutrition* 131: 1409–1413.
- Takeda K, Kaisho T, Akira S, 2003. Toll-like receptors. *Annual Review of Immunology* 21: 335–376.
- Talpur NA, Echard BW, Fan AY, Jaffari O, Bagchi D, Preuss HG, 2002. Antihypertensive and metabolic effects of whole maitake mushroom powder and its fractions in two rat strains. *Molecular and Cellular Biochemistry* 237: 129–136.
- Tam SC, Yip KP, Fung KP, Chang ST, 1986. Hypotensive and renal effects of an extract of the edible mushroom *Pleurotus sajor-caju*. *Life Science* 38: 1155–1161.
- Taylor PR, Tsoni SV, Willment JA, Dennehy KM, Rosas M, Findon H, Haynes K, Steele C, Botto M, Gordon S, Brown GD, 2007. Dectin-1 is required for β-glucan recognition and control of fungal infection. *Nature Immunology* 8: 31–38.
- Thornton BP, Vetvicka V, Pitman M, Goldman RC, Ross GD, 1996. Analysis of the sugar specificity and molecular location of the β-glucan-binding lectin site of complement receptor type 3 (CD11b/CD18). *Journal of Immunology* 156: 1235–1246.
- Tinchieri G, 2003. Interleukin-12 and the regulation of innate resistance and adaptive immunity. *Nature Reviews Immunology* 3: 133–146.
- Tsikitis VL, Morin NA, Harrington EO, Albina JE, Reichner JS, 2004. The lectin-like domain of complement receptor 3 protects endothelial barrier function from activated neutrophils. *Journal of Immunology* 173: 1284–1291.
- Underhill DM, Rossnagle E, Lowell CA, Simmons RM, 2005. Dectin-1 activates Syk tyrosine kinase in a dynamic subset

- of macrophages for reactive oxygen production. *Blood* 106: 2543-2550.
- Vassallo R, Standing JE, Limper AH, 2000. Isolated pneumocystis carinii cell wall glucan provokes lower respiratory tract inflammatory responses. *Journal of Immunology* 164: 3755-3763.
- Vetvicka V, Thornton BP, Ross GD, 1996. Soluble  $\beta$ -glucan polysaccharide binding to the lectin site of neutrophil or natural killer cell complement receptor type 3 (CD11b/CD18) generates a primed state of the receptor capable of mediating cytotoxicity of iC3b-opsonized target cells. *Journal of Clinical Investigation* 98: 50-61.
- Vetvicka V, Yvin JC, 2004. Effects of marine  $\beta$ -1,3 glucan on immune reactions. *International Immunopharmacology* 4: 721-730.
- Viriyakosol S, Fierer J, Brown GD, Kirkland TN, 2005. Innate immunity to the pathogenic fungus *Coccidioides posadasii* is dependent on toll-like receptor 2 and Dectin-1. *Infection and Immunity* 73: 1553-1560.
- Walsh TJ, Tepler H, Donowitz GR, Maertens JA, Baden LR, Dmoszynska A, Cornely OA, Bourque MR, Lupinacci RJ, Sable CA, dePauw BE, 2004. Caspofungin versus liposomal amphotericin B for empirical antifungal therapy in patients with persistent fever and neutropenia. *New England Journal of Medicine* 351: 1391-1402.
- Wang HX, Ooi VE, Ng TB, Chiu KW, Chang ST, 1996. Hypotensive and vasorelaxing activities of a lectin from the edible mushroom *Tricholoma mongolicum*. *Pharmacology and Toxicology* 79: 318-323.
- Wang J, Gigliotti F, Maggirwar S, Johnston C, Finkelstein JN, Wright TW, 2005. *Pneumocystis carinii* activates the NF-kappaB signaling pathway in alveolar epithelial cells. *Infection and Immunity* 73: 2766-2777.
- Wang Y, Zhang L, Li Y, Hou X, Zeng F, 2004. Correlation of structure to antitumor activities of five derivatives of a  $\beta$ -glucan from *Poria cocos sclerotium*. *Carbohydrate Research* 339: 2567-2574.
- Wei D, Williams D, Browder W, 2002a. Activation of AP-1 and SP1 correlates with wound growth factor gene expression in glucan-treated human fibroblasts. *International Immunopharmacology* 2: 1163-1172.
- Wei D, Zhang L, Williams DL, Browder IW, 2002b. Glucan stimulates human dermal fibroblast collagen biosynthesis through a nuclear factor-1 dependent mechanism. *Wound Repair and Regeneration* 10: 161-168.
- Williams DL, Di Luzio NR, 1980. Glucan-induced modification of murine viral hepatitis. *Science* 208: 67-69.
- Willment JA, Gordon S, Brown GD, 2001. Characterization of the human  $\beta$ -glucan receptor and its alternatively spliced isoforms. *Journal of Biological Chemistry* 276: 43818-43823.
- Willment JA, Marshall AS, Reid DM, Williams DL, Wong SY, Gordon S, Brown GD, 2005. The human  $\beta$ -glucan receptor is widely expressed and functionally equivalent to murine Dectin-1 on primary cells. *European Journal of Immunology* 35: 1539-1547.
- Xia Y, Borland G, Huang J, Mizukami IF, Petty HR, Todd 3rd RF, Ross GD, 2002. Function of the lectin domain of Mac-1/complement receptor type 3 (CD11b/CD18) in regulating neutrophil adhesion. *Journal of Immunology* 169: 6417-6426.
- Xia Y, Ross GD, 1999. Generation of recombinant fragments of CD11b expressing the functional  $\beta$ -glucan-binding lectin site of CR3 (CD11b/CD18). *Journal of Immunology* 162: 7285-7293.
- Xiao Z, Trincado CA, Murtaugh MP, 2004. B-glucan enhancement of T cell IFN gamma response in swine. *Veterinary Immunology and Immunopathology* 102: 315-320.
- Yanaguchi M, Matsuura M, Kobayashi K, Sasaki H, Yajima T, Kuwata T, 2001. Lactoferrin protects against development of hepatitis caused by sensitization of Kupffer cells by lipopolysaccharide. *Clinical and Diagnostic Laboratory Immunology* 8: 1234-1239.
- Yan J, Allendorf DJ, Brandley B, 2005. Yeast whole glucan particle (WGP)  $\beta$ -glucan in conjunction with antitumor monoclonal antibodies to treat cancer. *Expert Opinion in Biological Therapy* 5: 691-702.
- Yokota K, Takashima A, Bergstresser PR, Arizumi K, 2001. Identification of a human homologue of the dendritic cell-associated C-type lectin-1, dectin-1. *Gene* 272: 51-60.
- Yoshioka S, Ohno N, Miura T, Adachi Y, Yadomae T, 1998. Immunotoxicity of soluble  $\beta$ -glucans induced by indomethacin treatment. *FEMS Immunology and Medical Microbiology* 21: 171-179.
- Young RE, Thompson RD, Larbi KY, La M, Roberts CE, Shapiro SD, Perretti M, Nourshargh S, 2004. Neutrophil elastase (NE)-deficient mice demonstrate a nonredundant role for NE in neutrophil migration, generation of proinflammatory mediators, and phagocytosis in response to zymosan particles in vivo. *Journal of Immunology* 172: 4493-4502.
- Zekovic DB, Kwiatkowski S, Vrvic MM, Jakovljevic D, Moran CA, 2005. Natural and modified (1 $\rightarrow$ 3)- $\beta$ -D-glucans in health promotion and disease alleviation. *Critical Reviews in Biotechnology* 25: 205-230.
- Zhang L, Li X, Xu X, Zeng F, 2005. Correlation between antitumor activity, molecular weight, and conformation of lentinan. *Carbohydrate Research* 340: 1515-1521.
- Zimmerman JW, Linderthuth J, Fish PA, Palace GP, Stevenson TT, DeMong DE, 1998. A novel carbohydrate-glycosphingolipid interaction between a  $\beta$ -(1-3)-glucan immunomodulator, PGG-glucan, and lactosylceramide of human leukocytes. *Journal of Biological Chemistry* 273: 22014-22020.